

Dynamic properties of isolated acetylcholine receptor protein: Kinetics of the binding of acetylcholine and Ca ions*

(relaxation kinetics/temperature jump/relaxation time analysis/conformational change)

EBERHARD NEUMANN[†] AND HAI WON CHANG[‡]

[†]Max-Planck-Institut für Biochemie, D-8033 Martinsried bei München, Germany; and [‡]Department of Neurology, College of Physicians and Surgeons, Columbia University, New York, N.Y. 10032

Communicated by David Nachmansohn, July 13, 1976

ABSTRACT Kinetic and thermodynamic constants for elementary steps associated with the interaction of acetylcholine (AcCh) and Ca with isolated AcCh receptor from *Torpedo californica* have been determined by chemical relaxation spectrometry. Murexide is used as a Ca indicator to monitor changes in Ca bound to the AcCh receptor. In the presence of AcCh this technique permits an indirect monitoring of AcCh binding, because the AcCh and the Ca binding reactions are competitively coupled. A temperature-jump perturbation in the Ca-murexide-AcCh receptor system induces a spectrum of relaxation processes characterized by at least three relaxation times: $\tau^1 = 5 (\pm 1)$ msec; $\tau^2 = 35 (\pm 5)$ msec; and $\tau^3 = 300 (\pm 30)$ msec. In the presence of AcCh, the Ca relaxation spectrum is altered in a characteristic way. A formalism is developed to describe the normal mode relaxation times of the coupled reaction system in terms of total concentrations of both AcCh and receptor binding sites. The analysis also allows one to determine the stoichiometry of the reactions involved or to estimate a molecular weight of the AcCh receptor. The kinetic data suggest that the reaction of AcCh with receptor proceeds in at least two steps. The rate constant of the association of AcCh with receptor was found to be $2.4 (\pm 0.5) \times 10^7 \text{ M}^{-1} \text{ sec}^{-1}$ at 23.5°, 0.1 M NaCl, 50 mM Tris-HCl, pH 8.5. Reaction schemes consistent with the present kinetic data are discussed in terms of a physicochemical model that accounts for the rapid transient conductivity changes in excitable membranes during nerve and muscle excitation.

The conductance changes in excitable membrane during nerve and muscle excitation are controlled by physicochemical membrane processes. At present, however, the molecular mechanism that underlies such changes is unknown. One may anticipate that several reaction steps are involved that are coupled with each other, possibly in a sequential form (1). Consequently, the time constants of the measured electrical changes reflect normal mode time constants of the whole regulatory system.

An experimental approach toward an understanding of the molecular mechanism of excitability is a study of elementary reactions involving components thought to be involved in the excitation process. One such component is the acetylcholine receptor protein. Association of the neural activator acetylcholine (AcCh) with receptor appears to initiate a sequence of reactions which ultimately result in local permeability changes in excitable membranes (for review see refs. 1 and 2).

Recent experiments with highly purified AcCh receptor isolated from electric organs of electric fish indicate that the binding of AcCh (3) or other neural activators (4) induces release of Ca bound to AcCh receptor. In contrast, binding of α -bungarotoxin, an inhibitor, causes an uptake of Ca ions on the receptor (3). These observations suggest that AcCh and

α -bungarotoxin cause different conformational shifts in the AcCh receptor.

Study of the elementary reactions involving AcCh and AcCh receptor is complicated by the lack of a suitable reaction indicator. In this study these reactions are monitored indirectly through their coupling to Ca-AcCh receptor binding equilibria. With the use of murexide as an indicator of Ca ions, a spectrophotometric method has been developed to measure changes in the concentration of Ca ions bound to AcCh receptor (3). To the extent that AcCh reactions are coupled to the Ca binding, changes in bound Ca can be used to indicate changes caused by AcCh in presumed AcCh receptor conformational states. The reaction equilibria involving receptor, Ca, and AcCh are investigated here by the temperature-jump relaxation method, and kinetic and thermodynamic constants associated with elementary reaction equilibria are reported. The preliminary results of these studies have been briefly referred to in previous publications (1, 3, 5, 6).

MATERIALS AND METHODS

AcCh Receptor. The preparation of AcCh receptor used in the present study was freshly purified from electric tissue of *Torpedo californica* that had been frozen in liquid nitrogen, as described (3). The purified AcCh receptor samples had a binding capacity of 7.6–10 nM ¹²⁵I-labeled α -bungarotoxin per mg of protein. The protein preparations were characterized by a major AcCh binding component with an equilibrium dissociation constant $K_A = 1.5 (\pm 0.2) \times 10^{-6}$ M and a maximum of 2 AcCh molecules bound to 260,000 molecular weight (3). Immediately before the experiments, solutions of AcCh receptor in 0.1% Brij, 15 mM sodium phosphate buffer, 0.1 M NaCl, pH 7.0 were concentrated by vacuum dialysis in a collodion bag (Schleicher & Schuell) at 0° against three changes of buffer, 50 mM Tris-HCl, 0.1 M NaCl, pH 8.5 at 20° (buffer I) and centrifuged to remove particulate matter. The temperature-jump samples, 0.8 ml of solution, in buffer I, contained 2.4 mg of AcCh receptor and 80 μ M murexide. The stock solution of murexide (0.4 mM) in buffer I had been twice filtered through Millipore filters (0.45 μ m) before it was mixed with the AcCh receptor solutions. CaCl₂ and AcCh were added where appropriate from stock 50 mM and 5 mM solutions in buffer I, respectively, in Aglar microsyringes.

Kinetic Method. A Joule-heating temperature-jump spectrophotometer apparatus (7, 8) was used to measure the chemical relaxation spectrum of AcCh receptor. The apparatus, a modified version of a previously reported instrument (8), was further developed by C. R. Rabl, Göttingen, and was equipped with an electronic cooling correction device. The tempera-

Abbreviation: AcCh, acetylcholine.

* This paper is part II of a series. Paper I is ref. 3.




FIG. 1. Chemical relaxation spectrum (section $t \geq 5$ msec) of AcCh receptor from *T. californica*, at 1 mM Ca, 23.5°, after a temperature jump of 3.5°. ΔI measures light intensity changes at 480 nm. 10 mV correspond to an absorbance change of 0.0005 per 0.7 cm. Double-sweep oscilloscope record at two time scales; B₁ and B₂ are reference tracings (no temperature jump). The pre-amplitude of 578 mV ($t \leq 5$ msec) was electronically suppressed, and cooling correction was applied.

ture-jump cell contained 0.7 ml of carefully degassed temperature-jump samples described above. The electrical resistance, R , of the filled cell was 220 Ω . The capacitance, C , of the high voltage discharge capacitor was 5×10^{-8} F. The temperature increase per jump was between 3 and 5°. Under the given experimental conditions, the heating time $\tau_h \approx 0.5$ RC was 6 μ sec. The relaxation of the Ca-AcCh receptor system in the presence of murexide was monitored at 480 nm and 23.5° (see ref. 3).

Analysis. A relaxation time simulator was used to estimate relaxation amplitudes and time constants of the measured relaxation spectrum (10). A method was developed to analyze the estimated relaxation time constants in terms of the total concentrations of the reaction partners (11).

RESULTS

Ca-AcCh receptor relaxations

The concentrations of the reaction partners were chosen so that the equilibration of the indicator system Ca plus murexide is very rapid (relaxation time $\tau_{CaMu} = 10$ μ sec) and practically falls in the time range of the Joule heating of the measuring cell ($\tau_h = 6$ μ sec). The indicator system can then be used to monitor any slower relaxations involving Ca ions (7, 9). Typical relaxation curves of isolated AcCh receptor from *Torpedo californica* in 1 mM Ca are shown in Fig. 1. Due to photoreaction of murexide and photomultiplier drift at the extremely high sensitivity of the recording device, the base lines are inclined as shown. For the time range $t \geq 0.1$ msec in 1 mM Ca, it was found that the Ca relaxation spectrum is characterized by at least three major relaxation times: $\tau^1 = 5 (\pm 1)$ msec, $\tau^2 = 35 (\pm 5)$ msec, and $\tau^3 = 300 (\pm 30)$ msec (mean deviation in parentheses).

The relaxation spectrum of Ca-AcCh receptor varies with changes in the Ca concentration. However, the Ca concentration dependence of both time constants and relaxation amplitudes is small and rather complicated. The present accuracy of the data is not sufficient to differentiate between alternative mechanisms for the Ca-AcCh receptor interactions. It appears unlikely, however, that the two larger relaxation times in particular reflect simple bimolecular equilibria between Ca ions and AcCh receptor binding sites.

Changes caused by AcCh

In the presence of AcCh, the relaxation spectrum of the Ca-AcCh receptor system changes. As shown in Fig. 2, the relaxation times τ^1 and τ^2 pass through a maximum as a function of




FIG. 2. Kinetic titration with AcCh of AcCh receptor from *T. californica*. Plot of time constants τ^i of the Ca-AcCh receptor relaxation at 1 mM Ca and 23.5° as a function of $q = [A^0]/[R^0]$, where $[A^0]$ is the (variable) total concentration of AcCh and $[R^0]$ is the (practically constant) total concentration of AcCh binding sites of AcCh receptor (5×10^{-6} M). See text.




AcCh concentration, whereas the large time constant τ^3 practically remains constant.

DISCUSSION

Ca Binding to AcCh Receptor. The chemical relaxation spectrum obtained for the Ca-AcCh receptor system reflects multiple interactions between Ca ions and receptor. This observation is consistent with the results of thermodynamic Ca binding studies, which indicate multiple Ca binding sites on the AcCh receptor (3). The kinetics of metal ion binding to many reaction partners is usually close to diffusion controlled, and the rate constant of association of Ca ions with simple ligands at 0.1 M ionic strength and 20° is usually in the order of 10^8 - 10^9 (M sec)⁻¹ (7). As seen in the next section, for $[Ca] = 1$ mM, bimolecular elementary reactions involving Ca relax in the 20 μ sec time range. Therefore, the values of τ^2 and τ^3 indicate that slower intramolecular reactions, such as conformational changes, are reflected in the Ca-AcCh receptor relaxation.

Kinetics of AcCh Binding to Receptor. An analysis of the relaxation times represented in Fig. 2 suggests the presence of conformational equilibria of AcCh receptor. The isolated AcCh receptor equilibrates between at least two conformational states $R \rightleftharpoons R''$, having different apparent equilibrium constants for AcCh and Ca ions.

As shown in our previous report (3), the binding sites for Ca and AcCh may be treated as indistinguishable and independent. Furthermore, the dependence of the AcCh binding constants on the Ca concentration suggests competition between AcCh and Ca for the AcCh receptor macromolecule (3). In the simplest case, if such competition involves binding sites in AcCh receptor, R, to which both Ca and AcCh ions can bind, then a minimum reaction scheme that accounts for the observed dependence on AcCh, A, of the observed relaxations must consist of at least three elementary steps:



The experimental data show that $[R'']$ is very small, so that the bimolecular step $A + R'' \rightleftharpoons AR''$ is not visible.

In the above scheme, the k terms are the rate constants of the elementary steps associated with the equilibrium constants $K_{Ca} = k_{10}/k_{01}$, $K_1 = k_{21}/k_{12}$, and $K_2 = k_{23}/k_{32}$, respectively. The kinetic data suggest that the observed equilibrium constant K_A for the interaction of AcCh with receptor, determined by thermodynamic methods (equilibrium binding of $[^3H]AcCh$), is actually an overall equilibrium constant. As given in the reaction scheme, the reaction partners for AcCh are R and R'' . Therefore,

$$K_A = \frac{[A]([R] + [CaR])}{[AR] + [AR'']} = (K_1 \phi^{-1}) / (1 + K_2) \quad [4]$$

where $\phi = (1 + K_{Ca}^{-1}[Ca])^{-1}$. As shown previously (3), the dependence of K_A on $[Ca]$ is associated with $K_{Ca} = 3.1 (\pm 0.2) \times 10^{-3}$ M. Thus, at $[Ca] = 1$ mM, $\phi = 0.76$. It is now noted that Eq. 6 in ref. 3 must be specified to $K_A = K_1(1 + K_2)^{-1}(1 + K_{Ca}^{-1}[Ca])$.

Due to the coupling of the elementary reactions, the measured relaxation times are normal mode time constants of the reaction scheme given above. As has already been discussed, if $k_{01} = 10^8$ (M sec) $^{-1}$, then for the present conditions the intrinsic relaxation time for Eq. 1, given by $\tau_0 = \{k_{01}([Ca] + [R] + K_{Ca})\}^{-1}$ is calculated to be about 2.5 μ sec. The bimolecular Ca-AcCh receptor relaxation can thus safely be considered to be very rapid compared to the measured (and kinetically resolvable) time constants (see Fig. 2). It will be seen that the dependence of τ^1 on AcCh concentration suggests that this time constant is the normal mode relaxation time, τ_1 , associated with the coupled reaction in Eq. 2. Since $\tau_0 \ll \tau_1$, the relaxation time $\tau^1 = \tau_1$ includes the equilibration factor $\{1 + [Ca](K_{Ca} + [R])^{-1}\}^{-1}$ which, for the present experimental condition of $K_{Ca} \gg [R]$, can be approximated to ϕ (see Eq. 4).

For the conditions of a rapid equilibration in Eq. [1],

$$1/\tau_1 = k_{12}\phi([A] + [R]\phi^{-1}) + k_{21} = k_{12}' + k_{21} \quad [5]$$

Applying a procedure recently suggested by Eigen and Winkler-Oswatitsch (11), it is possible to express τ_1 in terms of total concentrations of AcCh, $[A^0]$, and of total binding sites of AcCh receptor, $[R^0]$. Using the definitions $q = [A^0]/[R^0]$ and $p = K_A/[R^0]$ and introducing $[A] = [A^0] \cdot ([AR] + [AR''])$ and $([R] + [CaR]) = [R^0] \cdot (1 + K_{Ca}^{-1}[Ca]) = [R^0] \cdot ([AR] + [AR''])$ into Eq. 4, it can be written that $2([AR] + [AR'']) = [R^0] \{ (1 + p + q) - \sqrt{(1 + p + q)^2 - 4q} \}$. Now, Eq. 5 can be rewritten to:

$$1/\tau_1 = k_{12}\phi[R^0] \left(\sqrt{(1 + p + q)^2 - 4q} + p \cdot K_2 \right) \quad [6]$$

If $p < 1$, i.e., $[R^0] > K_A$, τ_1 passes through a maximum as a function of q , at $q^m = 1 - p$.

The dependence of τ^2 on q suggests that this relaxation is associated with the intramolecular reaction in Eq. 3. With $\tau_{11} = \tau^2$ and since $\tau_{11} > \tau_1$, the reaction represented in Eq. 2 may be considered to be equilibrated during the slower relaxation of the intramolecular step. Introducing the equilibration factor $k_{12}'/(k_{12}' + k_{21})$, the relaxation time of the slow mode is given by

$$1/\tau_{11} = k_{23} \{ k_{12}' / (k_{12}' + k_{21}) \} + k_{32} \quad [7]$$

See refs. 7 and 11. With the definitions of p and q , Eq. 7 is rewritten as:

$$\tau_{11} - \tau_2 = \tau_2 \cdot \frac{p \cdot K_2}{\sqrt{(1 + p + q)^2 - 4q}} \quad [8]$$




FIG. 3. The ratio $r_{11} = (\tau_{11} - \tau_2) / (\tau_{11}^m - \tau_2)$ as a function of q , plotted according to Eq. 11. See text.

where

$$\tau_2 = (k_{21} + k_{12})^{-1} = \{k_{12}(1 - K_2^{-1})\}^{-1} \quad [9]$$

is the intrinsic time constant of the uncoupled reaction represented by Eq. 3.

If $p < 1$, τ_{11} passes through a maximum as a function of q at $q^m = 1 - p$. The time constant τ_{11}^m of the maximum is given by

$$\tau_{11}^m - \tau_2 = (1/2)\tau_2 K_2 \sqrt{p} \quad [10]$$

The combination of Eqs. 8 and 10 results in an expression for the ratio, r_{11} ,

$$r_{11} = \frac{\tau_{11} - \tau_2}{\tau_{11}^m - \tau_2} = \frac{2\sqrt{p}}{\sqrt{(1 + p + q)^2 - 4q}} \quad [11]$$

where the measurable relaxation times τ_{11} and τ_2 (which equals τ_{11} at $q \gg 1$) are only correlated with the variables q and p . The shape of the plot of r_{11} against q gives the value of p and thereby of K_A .

Recall that τ_1 and τ_{11} pass through a maximum as a function of q . In some preparations of AcCh receptor the position of the maximum corresponded to q values lower than $1 - p$; in these cases only a fraction of the protein appeared to bind AcCh.

In Fig. 3 the measured relaxation times τ_{11} are replotted according to Eq. 11 and compared with the calculated function $r_{11} = f(q, p)$ using $p = 0.2$ and $\tau_2 = 20$ msec. With $[R^0] = 5 \times 10^{-6}$ M on the basis of a molecular weight of 260,000 (see ref. 3), the kinetic method gives $K_A = p/[R^0] = 1.0 (\pm 0.5) \times 10^{-6}$ M, in good agreement with $K_A = 1.5 (\pm 0.3) \times 10^{-6}$ M derived from equilibrium dialysis binding data with this AcCh receptor preparation. When the maximum value $\tau_{11}^m = 50$ msec is inserted in Eq. 10, $K_2 = 6.7$; i.e., about 90% of complexed AcCh receptor is in the conformation AR'' , under the present experimental conditions. Inserting the K_2 value in Eq. 4, the calculation yields $K_1 = 6 \times 10^{-6}$ M. Thus, the kinetic method provides the individual equilibrium constants for the single steps whereas thermodynamic procedures only give the overall distribution constant, K_A . With Eq. 9 and $K_2 = 6.7$, the rate constants for the conformational change are $k_{23} = 43.5$ sec $^{-1}$ and $k_{32} = 6.5$ sec $^{-1}$.

The value of K_2 is now used to analyze the dependence on q of the relaxation time $\tau_1 = \tau^1$ (see Fig. 2). It is found that, for $k_{12} = 2.4 (\pm 0.5) \times 10^7$ (M sec) $^{-1}$, the measured time constants agree within experimental accuracy with the curve calculated according to Eq. 6. Since $k_{21} = K_1 \cdot k_{12}$, we find $k_{21} = 144 (\pm 30)$ sec $^{-1}$. Thus, the individual rate coefficients and equilibrium constants of the interaction between AcCh and receptor are obtained from relaxation data of Ca-receptor interactions.

11. Eigen, M. & Winkler-Oswatitsch, R. (1976) *Angew. Chem.* **88**, in press.
12. Gutfreund, H. (1972) in *Enzymes: Physical Principles* (Wiley-Interscience, London), p. 159.
13. Magleby, K. L. & Stevens, C. F. (1972) *J. Physiol. (London)* **223**, 173-197.
14. Katz, B. & Thesleff, S. (1957) *J. Physiol. (London)* **138**, 63-80.
15. Eldefrawi, M. E., Eldefrawi, A. T. & Wilson, D. B. (1975) *Biochemistry* **14**, 4304-4310.
16. O'Brien, R. D. & Gibson, R. E. (1975) *Arch. Biochem. Biophys.* **169**, 458-463.

Pr
V
Bi

N
a
b

D
*
Sal
Ce

Al
ph
pr
wi
me
ge
ha
Ge
in
reg
is
ir
the
clu
po
in
lip
—
Ur
3-
in
a
c

i.e.
wil
pro
shc
DF
bet
(so
fir
noi
ad
per
per
tha
(fr
siti
son
(2
tra
] son
29
line
the
anc
—
Abt
cho
NM