

## The *sim* Gene of *Escherichia coli* Phage P1: Nucleotide Sequence and Purification of the Processed Protein

JULIA MAILLOU AND BRIGITTE DREISEIKELMANN<sup>1</sup>

Universität Bielefeld, Fakultät für Biologie, Lehrstuhl für Gentechnologie/Mikrobiologie,  
Postfach 8640, 4800 Bielefeld, Federal Republic of Germany

Received October 12, 1989; accepted November 2, 1989

The *sim* gene of bacteriophage P1 causes exclusion of a superinfecting P1 phage. We determined the nucleotide sequence of a 1.9-kb DNA fragment that, in plasmids, causes Sim phenotype. There are two open reading frames within this region for proteins of 82 and 259 amino acids. A 1.3-kb fragment containing the larger open reading frame was inserted into an expression vector. Induced cells carrying the hybrid plasmid, termed pBD5, were not infected by phage P1 and produced a 24-kDa protein and, to a smaller extent, a 25-kDa protein. The 24-kDa protein was purified. Comparison of its amino-terminal amino acid sequence with the nucleotide sequence indicated that it is processed from a precursor protein by removal of a hydrophobic leader peptide of 20 amino acids. *In vivo* processing depends on *secA* gene function and is necessary for Sim interference with P1 infection. The data are discussed with respect to the function of the *sim* gene in superinfection exclusion. © 1990 Academic Press, Inc.

### INTRODUCTION

The *sim* gene of bacteriophage P1 was detected during attempts to clone the *c1* repressor gene (Devlin *et al.*, 1982). Carried by a multicopy vector, the *sim* gene confers immunity to cells against infection with wild-type P1 and also *c1*, *c4*, and *vir* mutants. Therefore the extended immunity was called superimmunity (*sim*). The *sim* gene is localized on the P1 *EcoRI*-9 DNA fragment within the *imm1* region (Bächi and Arber, 1977; Yarmolinsky, 1987).

In a previous study (Kliem and Dreiseikelmann, 1989), we showed that cells carrying a multicopy plasmid with the *sim* gene (pMK4) are not really immune. Such cells are not lysogenized, can adsorb phage, but fail to synthesize phage-specific proteins following infection. The *sim* gene product does not interfere with the circularization and replication of phage P1 DNA or with the maturation or release of progeny phages. We concluded that the *sim* gene product blocks some early step following infection but not transformation, probably at a stage between adsorption of the phage and circularization of the injected DNA. Thus the *sim* gene does not confer immunity but rather a superinfection exclusion (Kliem and Dreiseikelmann, 1989). Genes which may have a similar function have been described in other bacteriophages, such as *sieA* in phage P22 (Susskind *et al.*, 1974) and *immT* in phage T4 (Anderson and Eigner, 1971). The actual mechanism for superinfection exclusion is not yet known. The

genes have not been sequenced and the gene products have not been purified or characterized. It has been speculated that the *immT* gene product may be a nuclease or may activate a periplasmic endonuclease of *E. coli* (Anderson and Eigner, 1971).

Plasmids with the cloned *sim* region of P1 express the Sim phenotype and produce three proteins in a minicell system (Kliem and Dreiseikelmann, 1989). We have now sequenced the *sim* region and have identified and purified a 24-kDa protein which seems to be the protein responsible for the superinfection exclusion.

### MATERIALS AND METHODS

#### Bacterial strains

*Escherichia coli* JM101  $\Delta lac pro, thi, supE44$  [*F' traD36, proAB, lacI<sup>q</sup>Z M15*] was used for transformation with pUC13 and M13mp8 and M13mp9 RF derivatives (Yanisch-Perron *et al.*, 1985). *E. coli* C600  $F^-$ , *thi-1, thr-1, leuB61, lacY1, supE44, tonA21* was used for transformation with pJF118EH. Expression of genes from pPLc2819 derivatives was performed in *E. coli* K12  $\Delta H1 \Delta trp, Sm^r, lacZ, \Delta bio-uvrB, \Delta trpEA2$ , ( $\lambda$  Nam7-Nam53cl857  $\Delta H1$ ) (Remaut *et al.*, 1981). *E. coli* MM52  $F^-$ ,  $\Delta (lac)U169, araD136, relA, rspL, thi, secA51_{ts}$  was a gift from J. Tommassen (Oliver and Beckwith, 1981).

#### DNA sequencing

The nucleotide sequence was determined with the chain termination method (Sanger *et al.*, 1977). Various

<sup>1</sup> To whom requests for reprints should be addressed.



Fig. 1. Strategy for determination of the nucleotide sequence of a 1.9-kb DNA fragment from the *sim* region. Sequence of the upper strand was mainly determined with the aid of deletion derivatives produced by limited digestion with exonuclease III and VII (Yanisch-Perron *et al.*, 1985). The lower strand was sequenced after subcloning of different DNA restriction fragments.

DNA inserts were ligated into M13mp8 and mp9 RF-DNA (Messing and Vieira, 1982). DNA synthesis at the M13 single strand was performed with a T7 DNA polymerase system purchased from U.S. Biochemical Corporation (Tabor and Richardson, 1987).

### SDS-PAGE

Polyacrylamide slab gels (14 or 17.5% acrylamide, 1 mm thick, 8 cm long) were prepared as described (Laemmli, 1970). Electrophoresis was performed for about 2 hr at 20 mA. Gels were stained with Coomassie brilliant blue G-250. The protein marker mix from Pharmacia contained phosphorylase b (mol wt 94,000), BSA (mol wt 67,000), ovalbumin (mol wt 43,000), carboxyanhydrase (mol wt 30,000), trypsin inhibitor (mol wt 20,100),  $\alpha$ -lactoalbumin (mol wt 14,400).

### Isolation of the 24-kDa protein

**Growth of the cells and induction.** K12  $\Delta$  H1  $\Delta$  *trp* (pBD5) was grown in 1 liter TBY medium at 30° to a density of  $3 \times 10^8$  cells/ml. Temperature was shifted to 42° for 20 min and incubation was continued for 2 hr at 37°. Cells were harvested by centrifugation. The cells (wet wt 4 g) were resuspended in 5 ml cooled buffer A (20 mM Tris-HCl, pH 7.8, 40 mM NaCl, 0.1 mM EDTA, 7 mM  $\beta$ -mercaptoethanol).

**Crude extract.** Cells were disrupted by a French Press at a pressure of 18,000 psi. The lysate was centrifuged for 1 hr at 30,000 rpm in a TST55.5 rotor (Kontron) at 4°.

**Streptomycin sulfate step.** Streptomycin sulfate (30% solution) was slowly added to the crude extract to a final concentration of 3%. The mixture was stirred for 30 min at 0° and centrifuged for 30 min at 8000 rpm in a A8.24 rotor (Kontron).

**Ammonium sulfate precipitation.** Solid ammonium sulfate was added to the supernatant to 60% saturation. The mixture was stirred in ice for 30 min. After 2 hr at 0° the precipitate was collected by centrifugation

for 30 min at 8000 rpm in a A8.24 rotor. The precipitate was resuspended in 4 ml buffer B (20 mM sodium phosphate, pH 7.8, 0.1 mM EDTA, 7 mM  $\beta$ -mercaptoethanol) and dialyzed for 2 hr against the same buffer.

**Hydroxyapatite chromatography.** The dialyzed fraction was loaded onto a hydroxyapatite column (diameter 1.5 cm, height 2.5 cm), previously equilibrated with buffer B. The column was washed until the OD<sub>280</sub> was 0.05. Adsorbed proteins were eluted with a linear gradient (100 ml) of 20 to 200 mM sodium phosphate in buffer B. The flow rate was 10 ml/hr. Fractions of 1 ml were collected. The 24-kDa protein was identified by SDS-PAGE. Fractions containing this protein, which eluted between 40 and 80 mM sodium phosphate, were pooled and dialyzed against buffer A.

**Mono Q ion-exchange chromatography.** The dialyzed sample was loaded on an equilibrated Mono Q column (HR5/5, Pharmacia) and chromatographed by FPLC. The 24-kDa protein was eluted at a NaCl concentration of 300 to 400 mM NaCl with a linear gradient from 0.05 to 1 M NaCl in buffer A.

**Protein sequencing.** The protein was transferred to a membrane (Immobilon, Millipore) and the amino-terminal amino acids were determined with an automated protein sequencer (Knauer model 810, Berlin). Separation of the PTH amino acids was performed on-line on a PTH-C<sub>18</sub> column (Applied Biosystems, 220  $\times$  2.1 mm) at a flow rate of 0.24 ml/min using a sodium acetate/THF/acetonitrile eluent system as described (Hunkapiller, 1985).

## RESULTS

### Nucleotide sequence analysis of the *sim* region

In a preceding paper (Kliem and Dreiseikelmann, 1989) we have shown that the *sim* gene product interferes with P1 infection by blocking an early step following phage adsorption. In minicells three proteins with apparent molecular weights of about 25, 24, and 15 kDa were detected originating from the 2.1-kb *EcoRI*-*PvuII* *sim* region. In order to sequence the gene(s) re-

```

      10      20      30      40      50      60
CCCCATCAGCCCCACCATAATTTCGGGCGTAACGCGGTTTACGGACACAAAAATACCGCA

      70      80      90      100     110     120
ATATCGGAAATCTGCGGTGTCCGCACTAACATTTCAGGCTGTCAAACCGGTGCGAGAAT
      ← c4 →

      130     140     150     160     170     180
TTGCTACGACGGTGGAACTATAAGCCTGAACGATTAAGGTCAATATGATGCGAAAAGA

      190     200     210     220     230     240
TAGCATTTCGCGACTTAAAAATACAAATTTATTAGAGCATTATTTGTTTAAATAATACACA

      250     260     270     280     290     300
GTTGGATCTAATAACCTCTTTTTTTTAAAGGCGAAAATATGTACCCTAAATGAGTTATAA

      310     320     330     340     350     360
GGCAGGTGAGGTTATAATGAGAAAACCTACTACCGTTATTATTTATGGCTGGGACTGT

      370     380     390     400     410     420
TAATGCAGCATCAAGCGTAAAGGAGATTTGTACCGATTATACGAAATACCTTGGGCACGT

      430     440     450     460     470     480
TTACGCCCTTTGCGATCAATGACTATTAATCCAGTGCTAGGAGTGGATTTATGCTGATTCG
      -35          -10          SD          MetLeuIleAr

      490     500     510     520     530     540
TTTGTTTTTAGTGCTTTCCTTTTTAACATTTAATGTTTTGCTGATGAAGTTGACTTTTC
gLeuPheLeuValLeuSerPheLeuThrPheAsnValPheAlaAspGluValAspPheSe

      550     560     570     580     590     600
GAAGGTAGATTGCAATTCAGTGGAAACAAGAAAAGCTCTTATTGAAGAATATAACGAAAT
rLysValAspCysAsnSerValGluThrArgLysAlaLeuIleGluGluTyrAsnGluIl

      610     620     630     640     650     660
ATTATCGTCATATGGAATAACAGTGGTTGATTCTTATAATCAAAAAACTATTCAGAAAGG
eLeuSerSerTyrGlyIleThrValValAspSerTyrAsnGlnLysThrIleGlnLysGl

      670     680     690     700     710     720
AATAAATAAACTGGTCTGTTATGGGGTTTACCAATATTCAGATGGCTCTTCGGAGTAGTT
yIleAsnLysLeuValCysTyrGlyValTyrGlnTyrSerAspGlySerSerGlu

      730     740     750     760     770     780
TATTTATAAGCATAACCCAAATGTCTGTGAATTAATTAGTGAGTTTAAACCGATTAATGAG

```

FIG. 2. Nucleotide sequence of a 1.9-kb DNA fragment of the *sim* region. The two open reading frames which could encode proteins are shown with their deduced amino acid sequences. A possible promoter and the Shine–Dalgarno (SD) sequences are marked.

sponsible for the superinfection exclusion phenotype within this region, we inserted various subfragments of the *EcoRI*–*PvuII* fragment into M13mp8 and mp9 RF-DNA (Bächi and Arber, 1977). The sequencing strategy is summarized in Fig. 1. The nucleotide sequence determined by dideoxy sequencing is shown in Fig. 2.

At the 5' end of the nucleotide sequence there is part of the *c4* gene reading from right to left (position 93–1). The sequence agrees with published data (Baumstark and Scott, 1987). Two other open reading frames with the opposite orientation as the *c4* gene (positions 1524–1381 and 1272–1132), and one with the same

orientation (position 290–445), are evident. However these lack apparent transcriptional (promoter) and translational (ribosome binding site) signals. Two open reading frames which could possibly encode proteins are indicated along with their amino acid sequences in Fig. 2. One open reading frame starts at position 470 and ends at position 715. Upstream of it there is a possible promoter with a typical –10 and –35 sequence and a Shine–Dalgarno consensus sequence. The open reading frame would encode a protein of 82 amino acids. The second open reading frame (position 776–1547) has two start codons (position 776 and 791) but

790            800            810            820            830            840  
 TGAATGAAAATGAAATTATTTAATGTAATAACATTTTGTGTGCTATTTTTGCTGGAAG  
 5D            MetLysLeuPheAsnValIleThrPheCysCysAlaIlePheAlaGlySe

850            860            870            880            890            900  
 CGCGATAGCTGATAATAAATTGCCAGATTGGCTTTCTACCTCAAAGAAAGATTATGATTT  
 rAlaIleAlaAspAsnLysLeuProAspTrpLeuSerThrSerLysLysAspTyrAspLe

910            920            930            940            950            960  
 AGTAAGGGCATTCTATTTGTCTGGATTGCTTCGAAAGCAATGAACAATCAATTTGGTTA  
 uValArgAlaPheTyrLeuSerGlyPheAlaSerLysAlaMetAsnAsnGlnPheGlyTy

970            980            990            1000            1010            1020  
 TCATTTGCCATCTGAGTTGGTTAATGATTTTAAAGATAATGAATTTGCTGCTCAGAAAA  
 rHisLeuProSerGluLeuValAsnAspPheLysAspAsnGluPheAlaAlaGlnGluLy

1030            1040            1050            1060            1070            1080  
 ATGGAACACAATTCCAATTGTGTATGGTGAATAAAATCCATAAGAATGGTGAATAATAA  
 sTrpAsnThrIleProIleValTyrGlyGluIleLysSerIleArgMetValAsnAsnLy

1090            1100            1110            1120            1130            1140  
 ACCAATTGTAGAATTATTTACTCCAGGGGAAAATGCAACGCCTATAAATTATATCAAATT  
 sProIleValGluLeuPheThrProGlyGluAsnAlaThrProIleAsnTyrIleLysLe

1150            1160            1170            1180            1190            1200  
 GAAAATATTGGATTCAAAGCAAGACTCTCTGTTAAACTAAAAAAGGGGATGATATATA  
 uLysIleLeuAspSerLysGlnAspSerLeuLeuLysLeuLysLysGlyAspAspIleTy

1210            1220            1230            1240            1250            1260  
 TGCAGTGTGCTCCGGTGCTAATTTTAGCTTAGTGCCAATTCTGAGCAACTGCACTCCAGC  
 rAlaValCysSerGlyAlaAsnPheSerLeuValProIleLeuSerAsnCysThrProAl

1270            1280            1290            1300            1310            1320  
 AACAGACGTCATTGATGCTGCACTCTCTTTTCTGGTGAATATATGTTCCCTGCTTTTGA  
 aThrAspValIleAspAlaAlaLeuSerPheSerGlyGluTyrMetPheProAlaPheAs

1330            1340            1350            1360            1370            1380  
 TTCTTTTTCACCTACTAAGCAAACGTCAAATATATATTCACAAATCAAGATCCTGTTCAP  
 pSerPheSerProThrLysGlnAsnValLysTyrIlePheThrAsnGlnAspProValGl

1390            1400            1410            1420            1430            1440  
 GATGATGAATTTTATAGGGTACCTATCCTTAGTCGATACAACGAAGGATAAGAATAAAAT  
 nMetMetAsnPheIleGlyTyrLeuSerLeuValAspThrThrLysAspLysAsnLysMe

1450            1460            1470            1480            1490            1500  
 GGATATGGTTTCGTAAGTGTACGCCTGGAAGCCGGAATGTTACACAATTCGTTGATGT  
 tAspMetValArgLysCysThrProTrpLysProGluCysSerGlnGlnPheValAspVa

1510            1520            1530            1540            1550            1560  
 AATGGAAGGATTTGATAGCATCATGTATAAATATGAGGGAGAATTTAAAACTACATAGA  
 lMetGluGlyPheAspSerIleMetTyrLysTyrGluGlyGluPheLysAsnTyrIleGl

1570            1580            1590            1600            1610            1620  
 ATTAAATAGTGTGTTGAAAGGATAGTCAATTTAAGTTTAAACACCAACGCCATTAAAGG  
 uLeuLys

1630            1640            1650            1660            1670            1680  
 GCTTTTATTGTTTTACTCAAAACAACCTGATTAGTGTATAACCATATCATATCCCTCATT

1690            1700            1710            1720            1730            1740  
 CCACCTACACTGATTACCCCCAGACAACAATATTCCTACTCAATGAACAAATGACTACTC

1750            1760            1770            1780            1790            1800  
 GTAGAAFCGGTTAACACACCAGATTCTACGAGTTTCAATGACACCACGACAATTACTCG

1810            1820            1830            1840            1850            1860  
 AAGACGCAAAAACCGCTTCACACCTTTGATTGCGGATGAACCGGCCTTACTGGAATCCC

1870            1880            1890            1900  
 TGCTAAGAAAAGCATTTGGGAACCTACCAGGATAGGGCGGG

FIG. 2--Continued




Fig. 3. Hydropathic plot of the hypothetical Sim protein. The hydropathic plot was performed as described at a span setting of nine residues (Kyte and Doolittle, 1982).

only the AUG at 791 has a possible ribosome binding site, allowing for a protein of 259 amino acids. A promoter is not evident. At the end of the open reading frame there is an inverted repeat followed by numerous thymine residues, which may represent a transcriptional terminator (position 1606–1634).

The possible protein encoded by the latter open reading frame would have a molecular weight of 29,326 Da and may represent the 25-kDa protein previously described (Kliem and Dreiseikermann, 1989). The apparent discrepancy between the molecular weight deduced from the nucleotide sequence and that determined by SDS-PAGE will be considered in the discussion. For simplicity we will maintain the 24- and 25-kDa protein nomenclature. A hydropathic plot shows that the 25-kDa protein resembles a precursor protein with a hydrophobic leader sequence (Fig. 3). The amino-terminal amino acid sequence has the characteristics of a hydrophobic signal sequence (Oliver, 1985): (i) the length of the hydrophobic region is 20 amino acids, (ii) the protein begins with a positively charged region (Lys 2, Asn 5), (iii) there are two helix destabilizing amino acids at the distal site of the hydrophobic amino acid sequence (Gly 16 and Ser 17), (iv) the hydrophobic leader ends with the characteristic Ala-X-Ala sequence (Ala 18–Ile 19–Ala 20). This suggests that the 24-kDa protein observed in micells (Kliem and Dreiseikermann, 1989) may originate by processing of the 25-kDa protein.

#### Isolation of the 24-kDa protein and determination of the amino-terminal amino acid sequence

The following experiments demonstrate that the protein predicted from the nucleotide sequence is indeed

synthesized *in vivo* as a precursor (25 kDa) and is subsequently processed by proteolytic removal of a leader peptide to give the mature protein (24 kDa).

The 1320- and 1300-bp *EcoRI*–*HindIII* DNA fragments from pMK4d5 and pMK4d6 (Kliem and Dreiseikermann, 1989) were inserted into the expression vector pPLc2819 (Remaut *et al.*, 1981). Cells carrying hybrid plasmid pBD4 or pBD5 were induced for 2 hr at 42°. Total protein from uninduced and induced cells was separated by SDS-PAGE. Overproduction of the 24- and the 25-kDa protein was observed (Fig. 4). Generally the 25-kDa protein was less abundant than the 24-kDa protein. The 24-kDa protein was purified from induced cells as described under Materials and Methods (Fig. 5). To eliminate minor contaminants from fractions obtained after chromatography on a Mono Q column these were further purified by electrophoresis on a preparative SDS polyacrylamide gel. The protein purified in this way gave a single band on a silver-stained SDS-PAGE loaded with 2  $\mu$ g of protein (data not shown).

The purified protein was subjected to automated protein sequencing. The sequence of the first 15 amino acids was identical to that predicted from the nucleotide sequence analysis, but started with the aspartic acid residue at nucleotide position 851 corresponding to Asp 21. From this we conclude that the 24-kDa protein is the processed form of a precursor protein. The cleavage site is between Ala 20 and Asp 21. Processing would remove a hydrophobic leader sequence of 20 amino acids (see Fig. 3).




FIG. 4. Overproduction of a 25- and 24-kDa protein in cells with plasmids pBD4 and pBD5. SDS polyacrylamide gel (14%) of total protein from uninduced and induced cells. 1, marker proteins; 2, pBD4 without induction; 3, pBD4 with thermoinduction; 4, pBD5 without induction; 5, pBD5 with thermoinduction.

**Processing of the 25-kDa protein is essential for its biological activity**

The 1300-bp DNA fragment containing the *sim* gene was inserted into the expression vector pJF118EH with the *tac* promoter (Fürste *et al.*, 1986). The hybrid plasmid, called pBD6, was introduced into *E. coli* MM52




FIG. 5. Purification of the 24-kDa protein. SDS polyacrylamide gel (17.5%) with samples from crude extract (lane 2), ammonium sulfate precipitation (lane 3), hydroxyapatite fractions (lane 4), Mono Q fractions (lane 5), marker proteins (lane 1).




FIG. 6. Processing of the *Sim* protein in *secA<sub>ts</sub>* cells. SDS polyacrylamide gel (14%) with total protein from *Escherichia coli secA<sub>ts</sub>* pBD6. Molecular weight standards (lane 1); 30°, induction with IPTG (lane 2); 30°, uninduced cells (lane 3); 42°, induction with IPTG (lane 4); 42°, uninduced cells (lane 5).

*secA<sub>ts</sub>*. At the nonpermissive temperature, the precursor forms of many membrane proteins and periplasmic proteins are accumulated in *secA<sub>ts</sub>* cells (Oliver and Beckwith, 1981). Cells with pBD6 were induced with 4 mM IPTG at 30 and 42°, and total protein was analyzed by SDS-PAGE (Fig. 6). After induction at 30° the 24-kDa protein is the predominant protein as compared to the 25-kDa precursor. However, at 42° the reverse is observed—the 25-kDa protein accumulates and the processed form can hardly be detected. Therefore we conclude that the 25-kDa protein precursor is converted into the 24-kDa mature protein by proteolytic cleavage of the leader sequence depending on the *secA* function.

Induced *secA<sub>ts</sub>* cells with pBD6 are not infected by P1 at the permissive temperature of 30°, but they are susceptible to the phage at the nonpermissive temperature of 42° (Table 1). Thus the *secA*-dependent processing of the *Sim* protein is necessary for its function in blocking phage infection.

TABLE 1  
INFECTION OF *E. coli secA<sub>ts</sub>* CELLS WITH PHAGE P1

Strain	Infective centers (ml)	
	30°	42°
MM52 pJF118	$3 \times 10^9$	$8 \times 10^{10}$
MM52 pBD7	$<10^2$	$5 \times 10^{10}$

### *In vitro* properties of the Sim protein

It has been suggested for phage T4 that the *immT* gene product may be a nuclease (Anderson and Eigner, 1971). The purified Sim protein (Mono Q fraction) had no detectable nuclease activity when tested with linear double-stranded DNA under a variety of conditions. For T4 it has recently been described (Obringer, 1988) that the *sp* gene product, which is encoded by a gene in the vicinity of the *immT* gene, directly interacts with the phage tail. Phage infectivity is abolished by destruction of components of the injection apparatus. Incubation of purified Sim protein with phage P1 (15 to 60 min at room temperature; 20 mM Tris-HCl, pH 7.5, 10 mM NaCl, 5 mM MgCl<sub>2</sub>, 5 mM CaCl<sub>2</sub>) did not reduce the efficiency of plating of the lysate. This result suggests that the purified protein does not directly interact with some tail protein to inhibit the contact of phage particles with the receptor or the induction of a membrane channel for the injection of the DNA. The negative result does not exclude such interactions *in vivo*, which may need other factors.

### DISCUSSION

The nucleotide sequence of the *sim* region contains two open reading frames. The smaller one is preceded by a typical promoter region and a Shine-Dalgarno sequence. Downstream of the smaller open reading frame a larger one follows with its own Shine-Dalgarno sequence. It is likely that both genes are transcribed from the same promoter as an operon. Plasmids pBD4 and pBD5 still express the *sim* phenotype, although the first open reading frame is deleted in these plasmids. This implies that the gene product of the small reading frame is not essential for the superinfection exclusion phenotype. However, deletion of a *KpnI*-*HindIII* DNA fragment from plasmid pBD5, which removes about one-fifth of the distal part of the gene, resulted in the loss of the *sim* phenotype (data not shown). The product of the larger reading frame must therefore be the Sim protein. Its overproduction and processing in cells with pBD5 suffices to exclude infecting P1 phages from the cell.

The mechanism by which the Sim protein abolishes infection of cells by phage P1 is still not clear. We can now add the observation that the Sim protein may be a periplasmic or membrane-associated protein to our earlier results showing that the Sim protein blocks an early stage of P1 infection (Kliem and Dreiseikelmann, 1989). The following facts support this hypothesis: (i) the Sim protein is synthesized as a precursor with a hydrophobic leader sequence of 20 amino acid residues, (ii) only the processed form of the protein is biologically active, (iii) processing of the precursor is *secA*

dependent, since overproduction of the precursor is reduced at the nonpermissive temperature compared to the mature form at the permissive temperature. This is also observed for several other periplasmic and membrane proteins which are synthesized at a reduced level in *secA* cells and accumulate the precursor forms (Liss and Oliver, 1986; Strauch *et al.*, 1986).

The Sim protein is a rather hydrophobic protein which may account for the discrepancy between the apparent molecular weight as determined by SDS-PAGE (25 kDa, sometimes up to 27 kDa depending on the choice of marker proteins) and the predicted molecular weight deduced from the nucleotide sequence (29.3 kDa). In the literature one can find contrasting statements on the mobility of hydrophobic proteins during electrophoresis. There are indications for lower (Garten *et al.*, 1975) as well as for higher mobilities (Heller, 1978) than expected after heating in detergent. The Sim protein, which was always incubated for 5 min at 100° before electrophoresis, had a higher mobility than expected.

Since the Sim protein was isolated from a soluble fraction (see Materials and Methods), we suggest that the major portion of the protein may be localized in the periplasmic space. However, we cannot exclude the possibility that part of the protein may also be localized in the membrane. If the Sim protein is a membrane protein, the membrane association would probably be a specific one, so that overproduction of the protein would lead to a saturation of potential membrane binding sites. Further studies employing isolated membranes and osmotic shock proteins from Sim-producing cells are necessary to decide the localization of the Sim protein and to elucidate the mechanism of superinfection exclusion.

### ACKNOWLEDGMENTS

We thank Prof. Dr. Tschesche and his co-workers for sequencing of the protein and U. Beigel for excellent technical assistance. J. Mailou was supported by a grant from the European Community.

### REFERENCES

- ANDERSON, C. W., and EIGNER, J. (1971). Breakdown and exclusion of superinfecting T-even bacteriophages in *Escherichia coli*. *J. Virol.* **8**, 869-886.
- BÄCHI, B., and ARBER, W. (1977). Physical mapping of *BglII*, *BamHI*, *EcoRI*, *HindIII* and *PstI* restriction fragments of bacteriophage P1 DNA. *Mol. Gen. Genet.* **153**, 311-324.
- BAUMSTARK, B., and SCOTT, J. R. (1987). The *c4* gene of phage P1. *Virology* **156**, 197-203.
- DEVLIN, B. H., BAUMSTARK, B. R., and SCOTT, J. R. (1982). Superimmunity: Characterization of a new gene in the immunity region of P1. *Virology* **120**, 360-375.
- FÜRSTE, J. P., PANSEGRAU, W., FRANK, R., BLÖCKER, H., SCHOLZ, P., BAGDASARIAN, M., and LANKA, E. (1986). Molecular cloning of the

- plasmid RP4 primase region in a multi-host-range *tacP* expression vector. *Gene* **48**, 119–131.
- GARTEN, W., HINDENNACH, I., and HENNING, U. (1975). The major proteins of the *Escherichia coli* outer cell envelope membrane: Characterization of proteins II\* and III, comparison of all proteins. *Eur. J. Biochem.* **59**, 215–221.
- HELLER, K. B. (1978). Apparent molecular weights of a heat-modifiable protein from the outer membrane of *Escherichia coli* in gels with different acrylamide concentrations. *J. Bacteriol.* **134**, 1181–1183.
- HUNKAPILLAR, M. W. (1985). In "User Bulletin Protein Sequencer," Applied Biosystems Issue No. 14.
- KLIEM, M., and DREISEIKELMANN, B. (1989). The superimmunity gene *sim* of bacteriophage P1 causes superinfection exclusion. *Virology* **171**, 350–355.
- KYTE, J., and DOOLITTLE, R. F. (1982). A simple method for displaying the hydrophobic character of a protein. *J. Mol. Biol.* **157**, 105–132.
- LAEMMLI, U. K. (1970). Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* **227**, 680–685.
- LISS, L. R., and OLIVER, D. B. (1986). Effects of *secA* mutations on the synthesis and secretion of proteins in *Escherichia coli*. Evidence for a major export system for cell envelope proteins. *J. Biol. Chem.* **261**, 2299–2303.
- MESSING, J., and VIEIRA, J. (1982). A new pair of M13 vectors for selection either DNA strand of double-digest restriction fragments. *Gene* **19**, 269–276.
- OBRINGER, J. W. (1988). The functions of the T4 immunity and spackle genes in genetic exclusion. *Genet. Res.* **52**, 81–90.
- OLIVER, D. B. (1985). Protein secretion in *Escherichia coli*. *Annu. Rev. Microbiol.* **39**, 615–648.
- OLIVER, D. B., and BECKWITH, J. (1981). *E. coli* mutant pleiotropically defective in the export of secreted proteins. *Cell* **25**, 765–772.
- REMAUT, E., STANSENS, P., and FIERS, W. (1981). Plasmid vectors for high-efficiency expression by the  $p_i$  promoter of coliphage lambda. *Gene* **15**, 81–93.
- SANGER, F., NICKLEN, S., and COULSON, A. R. (1977). DNA sequencing with chain-termination inhibitors. *Proc. Natl. Acad. Sci. USA* **74**, 5463–5467.
- STRAUCH, K. L., KUMAMOTO, I., and BECKWITH, J. (1986). Does *secA* mediate coupling between secretion and translation in *Escherichia coli*? *J. Bacteriol.* **166**, 505–512.
- SUSSKIND, M. M., BOTSTEIN, D., and WRIGHT, A. (1974). Superinfection exclusion by P22 prophage in lysogens of *Salmonella thyphimurium*. III. Failure of superinfecting DNA to enter *sieA*<sup>+</sup> lysogens. *Virology* **62**, 350–366.
- TABOR, S., and RICHARDSON, C. C. (1987). DNA sequence analysis with a modified bacteriophage T7 DNA polymerase. *Proc. Natl. Acad. Sci. USA* **84**, 4767–4771.
- YANISCH-PERRON, C., VIEIRA, J., and MESSING, J. (1985). Improved M13 phage cloning vectors and host strains: Nucleotide sequences of the M13mp18 and pUC19 vectors. *Gene* **33**, 103–119.
- YARMOLINSKY, M. (1987). Bacteriophage P1. In "Genetic Maps, 1987" (S. J. O'Brien, Ed.), Vol. 4, pp. 38–47. Cold Spring Harbor Laboratory, Cold Spring Harbor, NY.