

## A Remark on Normal Forms of Matrices

Vlastimil Dlab

*Université de Poitiers*

*Poitiers, France*

and

Claus Michael Ringel

*Universität Bielefeld*

*Bielefeld, West Germany*

Submitted by Hans Schneider

---

### INTRODUCTION

This remark answers the two problems raised in [2]. As in [4], we use the recent techniques of [3] and [5] of the representation theory of finite-dimensional algebras. It seems that these techniques provide methods of solution, as well as proper understanding, of such classification problems.

### 1. FIRST PROBLEM

The first problem of [2] asks for normal forms of  $2m \times 2n$  complex matrices with respect to  $\mathbb{H}$ -similarity. Here, two complex  $2m \times 2n$  matrices  $A, A'$  are said to be  $\mathbb{H}$ -similar if there exist formally quaternionic invertible (square) matrices  $P, Q$  such that  $QA = A'P$ , and a complex  $2m \times 2n$  matrix  $P$  is called *formally quaternionic* if each block in its natural partition into  $2 \times 2$  blocks has the form

$$\begin{pmatrix} a & -\bar{b} \\ b & \bar{a} \end{pmatrix} \quad \text{with } a, b \in \mathbb{C},$$

where  $\bar{c}$  denotes the complex conjugate of  $c \in \mathbb{C}$ .

In order to solve this problem, one may just follow the general procedure presented in [4] and illustrated there on the classification problem of  $2m \times 2n$  real matrices with respect to  $\mathbb{C}$ -similarity. There, the problem was reformulated as the classification of real linear maps between two complex vector spaces. Similarly, in the present problem, we are concerned with the



- (ii) the corresponding transposed  $2p \times 2(p+1)$  matrices ( $p=1,2,\dots$ ), and
- (iii)  $2p \times 2p$  matrices ( $p=1,2,\dots$ )

$$\begin{pmatrix} E_c & E_1 & & & \\ & E_c & E_1 & & \mathbf{0} \\ & & \ddots & \ddots & \\ \mathbf{0} & & & E_c & E_1 \\ & & & & E_c \end{pmatrix} \quad \text{with } c = a + bi, a \geq 0,$$

or

$$\begin{pmatrix} E & E_1 & & & \mathbf{0} \\ & E & E_1 & & \\ & & \ddots & \ddots & \\ \mathbf{0} & & & E & E_1 \\ & & & & E \end{pmatrix},$$

where

$$E_c = \begin{pmatrix} 1-c & 0 \\ 0 & 1+\bar{c} \end{pmatrix} \quad \text{for complex } c \quad \text{and} \quad E = \begin{pmatrix} i & 0 \\ 0 & i \end{pmatrix}.$$

*These matrices are  $\mathbb{H}$ -indecomposable (i.e. not  $\mathbb{H}$ -similar to a proper direct product of two matrices), and in the decomposition of a complex  $2m \times 2n$  matrix, they are determined (up to their order) uniquely.*

## 2. SECOND PROBLEM

The second problem asks for normal forms of  $4m \times 4n$  real matrices with respect to  $\mathbb{H}$ -similarity. Here, two real  $4m \times 4n$  real matrices  $A, A'$  are said to be  $\mathbb{H}$ -similar if there exist formally real-quaternionic invertible (square) matrices  $P, Q$  such that  $QA = A'P$ , and a real  $4m \times 4n$  matrix  $P$  is called *formally real-quaternionic* if each block of its natural partition into  $4 \times 4$



in  $\text{End}_{\mathbb{R}}(X)$ , with

$$X = \bigoplus_m R \quad \text{and} \quad m = n + 2,$$

is just the set of  $R$ -multiples of the identity, and thus isomorphic to  $R$  (see [1]). Consider now  $Y_{\mathbb{H}} = X_{\mathbb{R}} \otimes_{\mathbb{R}} \mathbb{H}_{\mathbb{H}}$ . The centralizer of  $1 \otimes i$  and  $1 \otimes j$  (where  $i, j$  denote the corresponding left multiplications on  $\mathbb{H}$ ) in  $\text{End}(Y_{\mathbb{H}})$  are the elements  $\varphi \otimes 1$  with  $\varphi \in \text{End}(X_{\mathbb{R}})$ ; thus the centralizer of  $1 \otimes i$ ,  $1 \otimes j$  and  $\alpha \otimes 1 + \beta \otimes i$  in  $\text{End}(Y_{\mathbb{H}})$  will be isomorphic to  $R$ . However, this centralizer is precisely the endomorphism ring  $\text{End}_{\mathbb{H}}(A)$  of the real  $4m \times 4m$  matrix  $A$  corresponding to the representation

$$Y_{\mathbb{H}} \xrightarrow{\begin{matrix} 1 \otimes 1 \\ 1 \otimes i \\ 1 \otimes j \\ (\alpha \otimes 1) + (\beta \otimes i) \end{matrix}} Y_{\mathbb{H}}$$

of  $\mathbb{H} \xrightarrow{\mathbb{H}^4} \mathbb{H}$ . ■

It follows from this theorem that a classification of real  $4n \times 4n$  matrices with respect to  $\mathbb{H}$ -similarity is impossible: it would lead, at the same time, to a classification of all finite-dimensional  $\mathbb{R}$ -algebras.

### 3. CONCLUSION

Note that the first “open problem” is in fact, as we have shown above, a special case of the situation considered in the same paper [2]. It may perhaps be proper to emphasize two different objectives in dealing with classification problems: One is to find normal forms for a given problem; the other, usually easier objective is to show that two given problems have the same normal forms (modulo various discrete series of forms). The main theorem of [2] is a result of the second type (whereas the above solution of the first problem is of the first type). Let us remark that in such a situation, no simple normal form of matrices need exist at all—the classification of the similarity classes of matrices over a division ring seems to be a very difficult problem. On the other hand, the normal forms of matrices of discrete dimension type can always be listed [5], even in the “wild” situation of the second problem.

## REFERENCES

- 1 S. Brenner, Decomposition properties of some small diagrams of modules, *Symposia Mathematica* 13:128–142 (1974).
- 2 D. Ž. Djoković, Classification of pairs of maps, *Linear Algebra and Appl.* 20:147–165 (1978).
- 3 V. Dlab and C. M. Ringel, Indecomposable representations of graphs and algebras, *Mem. Amer. Math. Soc.* 173 (1976).
- 4 V. Dlab and C. M. Ringel, Normal forms of real matrices with respect to complex similarity, *Linear Algebra and Appl.* 17:107–124 (1977).
- 5 C. M. Ringel, Representations of  $K$ -species and bimodules, *J. Algebra* 41:269–302 (1976).

*Received 10 October 1978; revised May 1979*