

Kinetic Energy Release and Position of Transition State during Intramolecular Aromatic Substitution in Ionized 1-Phenyl-1-(2-pyridyl)ethylenes¹


Ralf Schubert and Hans-Friedrich Grützmacher*

Contribution from the Fakultät für Chemie der Universität Bielefeld, D-4800 Bielefeld 1, West Germany. Received September 4, 1979


Abstract: The loss of substituents (X = H, F, CH₃, Cl, Br, I) from the molecular ions of ortho-substituted 1-phenyl-1-(2-pyridyl)ethylenes **1a-f** and of the isomeric 1-phenyl-1-(3-pyridyl)- and 1-phenyl-1-(4-pyridyl)ethylenes **2** and **3** has been investigated. Cyclic fragment ions are formed from the ortho-substituted 1-phenyl-1-(2-pyridyl)ethylene molecular ions by an intramolecular aromatic substitution reaction. The energetic requirements of this reaction have been studied in dependence from the dissociation energy of the C-X bond by measurements of the ionization energies, appearance energies, and kinetic energies released during the reaction. The activation energy ϵ_a^* of the process varies only slightly with the dissociation energy of the C-X bond cleaved during the reaction, whereas the enthalpy of reaction changes from positive (endothermic) to very negative (exothermic) values in the reaction series **1a-f**. Consequently the reverse activation energy ϵ_r^* ranges from small to very large values in this series. This trend in ϵ_r^* is not followed by the kinetic-energy release. A large kinetic-energy release and energy partitioning quotient $q = 0.7-1.0$ is only observed for endothermic or thermoneutral processes, while a small kinetic-energy release and $q \approx 0.2$ is associated with exothermic reactions in spite of a large ϵ_r^* . This behavior has been correlated to the position X_0^* of the transition state on the reaction coordinate according to Miller's quantification of the Hammond postulate. The release of ϵ_r^* as kinetic energy is only observed for reactions with "symmetrical" or "late" transition states ($X_0^* > 0.4$) while most of ϵ_r^* remains as internal energy in the products of reactions with "early" transition states ($X_0^* < 0.4$).

The investigations of metastable ions in a mass spectrometer have been shown to reveal many mechanistic details of unimolecular reactions of organic ions.² The possibility of determining accurately the kinetic energy, T ,³ released during unimolecular reactions, is especially important. T stems from the potential energy of the transition state and the excess energy, ϵ^* , freely fluctuating in the activated complex. The nonfixed excess energy of metastable ions is usually small, and, if there is an activation energy of the reversed reaction, ${}^0\epsilon_r^*$, the major part of T often comes from ${}^0\epsilon_r^*$.^{3,4} In this case, ${}^0\epsilon_r^*$ can be approximated by the experimental activation energy of the reversed reaction, ϵ_r^* , which is derived from the appearance energy of the reaction and the sum of the heats of formation of the reaction products and which includes a small amount of nonfixed energy, ϵ^* , due to the kinetic shift of the appearance energy. The amount of ϵ_r^* , which appears as kinetic energy T in the products, can be expressed by the energy partitioning quotient $q = T/\epsilon_r^*$. The value of q depends obviously on the details of the energy hypersurface of the reaction in the neighborhood of the transition state. This has been shown for the elimination of H₂ and other small molecules from even-electron organic ions.^{2,5,6} The prerequisite for a large kinetic energy release during these reactions is not only a large ${}^0\epsilon_r^*$, but also a certain geometry of the activated complex, which allows channeling of its potential energy into translational energy of the products during the movement of the reaction system along the reaction coordinate.^{5,6} Hence a determination of q and its variation with structural changes in the reaction system offers the possibility of

Scheme I



Scheme II



(1) Mechanism of Mass Spectrometric Fragmentation Reactions. 26. Part 25: Schubert, R.; Grützmacher, H. F. *Org. Mass Spectrom.* **1980**, *15*, 122.

(2) Williams, D. H. *Acc. Chem. Res.* **1977**, *10*, 280 and references cited therein.

(3) Cooks, R. G.; Beynon, J. H.; Caprioli, R. M.; Lester, G. R. "Metastable Ions"; Elsevier: Amsterdam, 1973.

(4) Cooks, R. G.; Bertrams, M.; Beynon, J. H.; Rennekamp, M. E.; Setser, D. W. *J. Am. Chem. Soc.* **1973**, *95*, 1732.

(5) (a) Christie, J. R.; Derrick, P. J.; Richard, G. J. *J. Chem. Soc., Faraday Trans. 2* **1978**, *74*, 304. (b) Rickard, G. J.; Cole, N. W.; Christie, J. R.; Derrick, P. J. *J. Am. Chem. Soc.* **1978**, *100*, 2804. (c) Cole, N. W.; Rickard, G. J.; Christie, J. R.; Derrick, P. J. *Org. Mass Spectrom.* **1979**, *14*, 337.

(6) (a) Krause, D. A.; Day, R. J.; Jorgensen, W. L.; Cooks, R. G. *Int. J. Mass Spectrom. Ion Phys.* **1978**, *27*, 227. (b) Day, R. J.; Krause, D. A.; Jorgensen, W. L.; Cooks, R. G. *Ibid.* **1979**, *30*, 83.

observing some properties of the transition states of rather complex organic reactions directly.

In this paper the loss of ortho substituents at the phenyl group from molecular ions of substituted 1-phenyl-1-(2-pyridyl)ethylenes (**1a-f**, Scheme I) is discussed. Formally, this fragmentation corresponds to simple bond cleavage in an odd-electron ion, a frequently observed mass-spectrometric reaction. Its mechanism,

Table I. Partial 70-eV Mass Spectra^a of 1-Phenyl-1-pyridylethylenes and Substituted Derivatives

<i>m/z</i>	1a	1b	1c	1d	1e	1f	2	3
M	15	30	52	1	<1	<1	65	84
M-1		70	73	6	3	<1		
M-2		3	16					
M-3		4	15					
M-4			9					
180 (M-X)	100	100	100	100	100	100	100	100
179	4	7	7	3	5	6	6	4
178	7	11	7	7	9	11	8	6
177	1	3	2	2	2	2	2	2
166							23	10
153	2	1	2	1	2	2	9	17
152	17	9	13	8	11	17	35	41
151	6	8	5	4	5	8	14	15
150	2	3	1	1	1	3	4	4
140							3	2
139							5	5
128	1	1	3	1	1	2	2	3
127	3	1	2	2	2	3	6	9
126	1	2	1	1	1	3	4	4
115							3	4
114							1	1
113							2	1
102	2	5	1	1	2	3	5	6
101	1	2	1	3	2	3	2	2
90	4	3	5	2	5	9	6	5
89.5	1	3	-	4	2	2	4	5
89	3	4	5	3	2	1	6	6
79	2	4	6	1	1	1	1	1
78	4	8	6	4	4	5	4	4
77	10	4	5	3	4	9	16	22
76	3	3	2	2	2	5	10	13
75	2	7	2	3	2	4	5	5
63	2	2	3	1	1	3	6	7
52	4	8	5	1	2	3	3	4
51	5	8	5	4	4	8	12	14
50	2	3	2	1	1	3	5	5

^a Intensities in % base peak; peaks indicated represent >80% total ion current.

however, is better described as an intramolecular aromatic substitution reaction.^{7,8} In contrast to some other fragmentations of odd-electron aromatic molecular ions,⁴ for which constant energy partitioning is observed, the energy partitioning quotient *q* varies drastically with the nature of the substituent lost. It will be shown that the variation of *q* can be related to the position of the transition state at the reaction coordinate expressed by X_0^* , a function given by Miller⁹ for a "quantitative Hammond Postulate".

Results and Discussion

Mass Spectra and Structure of *m/z* 180 Ions. The 70-eV mass spectra of 1,1-diphenylethylene, 1-phenyl-1-(2-pyridyl)ethylene (**1a**), 1-phenyl-1-(3-pyridyl)ethylene (**2**), and 1-phenyl-1-(4-pyridyl)ethylene (**3**) (Table I) contain only a few intense signals due to fragment ions. While loss of a methyl radical is the predominant reaction of 1,1-diphenylethylene molecular ions besides loss of H and H₂, its aza analogues **1a**, **2**, and **3** fragment predominantly by loss of H, followed by elimination of CH₂N, to ions *m/z* 180 and 152, respectively.

The predominance of [M - H]⁺ ions (*m/z* 180) as primary fragmentation products is most easily explained by the formation of stable cyclic ions b' and b'' from **2**⁺ and **3**⁺, respectively, and b or a from **1a**⁺ (Scheme II). The reduced intensity of the molecular ions and the increased intensity of the [M - H]⁺ ions in the mass spectrum of **1a** suggest that the formation of ions a,

Table II. MI Spectra of [M - X]⁺ Ions (*m/z* 180; 70 eV)

<i>m/z</i>	1a	1b	1c	1d	1f	2	3
179 ^a	(583)	(259)	(194)	(205)	(176)	(398)	(260)
178	39	40	40	41	43	35	36
177	1	1	1	1	1		
153	27	18	19	19	14	49	61
152	28	35	34	35	36	16	13
151	5	6	6	5	6		

^a Omitted from normalization.¹²

Table III. CA Spectra of [M - X]⁺ Ions (*m/z* 180; 70 eV)

<i>m/z</i>	1a	1b	1c	1f	2	3
179 ^a	(108)	(115)	(86)	(94)	(81)	(60)
178 ^a	(130)	(145)	(121)	(128)	(89)	(72)
177	26	28	26	27	24	21
176	6	6	6	6	5	5
164-166	4	6	4	4	3	2
152-153 ^a	(53)	(49)	(47)	(47)	(86)	(88)
151 ^a	(32)	(30)	(30)	(31)	(47)	(50)
150	14	12	13	13	20	24
137-139	6	6	6	6	8	8
125-128	10	10	10	6	12	12
112-114	3	3	2	3	4	3
99-101	8	7	8	9	6	6
86-89	3	3	2	3	3	3
75-77	11	12	12	13	10	9
62-63	3	3	5	4	4	4
50-51	5	6	6	6	3	3

^a Omitted from normalization.

which are unique for **1a**, occurs with special ease. This is corroborated by the mass spectra of derivatives substituted at the phenyl group of **1a**, which show large peaks due to loss of the substituent X from the molecular ions only for the ortho isomers **1b-f** (see Table I and Figure 1).

The loss of a hydrogen atom or the substituent X and formation of *m/z* 180 ions is the only reaction observed for metastable molecular ions of **1a-f**, being obviously the only low-energy reaction path of these ions. The structures of the ions [M - X]⁺ (*m/z* 180) in the mass spectra of **1a-f**, **2**, and **3** have been studied by their unimolecular and collisional induced decompositions in the field-free region between the magnetic and electrostatic analyzer (second field-free region) of a VG ZAB 2F mass spectrometer using the DADI technique.¹⁰

All metastable *m/z* 180 ions formed in the 70-eV mass spectra of **1a-f**, **2**, and **3** decompose by losses of H, H₂, HCN, CH₂N, and CH₃N to product ions *m/z* 179, 178, 153, 152, and 151, respectively¹¹ (Table II).

The intensity pattern of the product ions of *m/z* 180 ions are identical within the limits of error for the ortho-substituted derivatives **1b-f**, which are expected to fragment to ions a.¹³ The *m/z* 180 ions arising from **2** and **3** differ clearly by a much larger

(10) (a) Beynon, J. H.; Caprioli, R. M.; Ast, T. *Org. Mass Spectrom.* **1971**, *5*, 229. (b) Maurer, K. H.; Brunnée, C.; Kappus, G.; Habfast, K.; Schröder, U.; Schulze, P. XIX Annual Conference on Mass Spectrometry and Allied Topics, Atlanta, Ga., 1971.

(11) As can be seen from the abnormal peak shapes, the signal at *m/z* 179 in the MI spectra of *m/z* 180 ions from **1a**, **2**, and **3** is composite and contains an interference peak besides the peak of (180 - H) ions. This interference peak stems from a metastable transition 182* → 181* occurring in the first field-free region and corresponding to loss of H from ¹³C molecular ions of **1a**, **2**, and **3**, which is transmitted by the instrument.¹² Hence the intensity of the signal at *m/z* 179 has to be excluded from a comparison of the MI spectra.

(12) Schaldach, B.; Grützmaier, H. F. *Org. Mass Spectrom.*, **1980**, *15*, 166.

(13) The *m/z* 180 ions in the mass spectra of **1a-f** are formed with a wide range of internal energies. This follows from the energetics of the fragmentations. It is of interest to note that this has no large effect on the intensity distribution of the product ions in the MI spectra. However, the total ion current of the MI spectra relative to the ion current of the mother ion is much less for *m/z* 180 ions from **1a-c** than from **1d-f**. The latter ions must contain a larger amount of excess energy than the former ones, which have been formed with a larger kinetic-energy release.

(7) Schaldach, B.; Grützmaier, H. F. *Int. J. Mass Spectrom. Ion Phys.* **1979**, *31*, 257.

(8) (a) Ronayne, J.; Williams, D. H.; Bowie, J. J. *Am. Chem. Soc.* **1966**, *88*, 4980. (b) Kuschel, H.; Grützmaier, H. F. *Org. Mass Spectrom.* **1974**, *9*, 408.

(9) Miller, A. R. *J. Am. Chem. Soc.* **1978**, *100*, 1984.




Figure 1. The 70-eV mass spectra of (a) 1-phenyl-1-(2-pyridyl)ethylene (**1a**), (b) 1-phenyl-1-(4-pyridyl)ethylene (**3**), (c) 1-(2-chlorophenyl)-1-(2-pyridyl)ethylene (**1d**), (d) 1-(4-chlorophenyl)-1-(2-pyridyl)ethylene.

intensity of m/z 153 product ions, indicating a different structure (probably b' and b''). No clear decision can be made for m/z 180 ions formed in the 70-eV mass spectrum of **1a**, however, since the intensity pattern falls between that observed for **1b** and **1f** and **2** and **3**, respectively, but likely metastable m/z 180 ions from **1a** represent a mixture of structures a and b.

The CA spectra of the m/z 180 ions from the 70-eV mass spectra of **1a-f**, **2**, and **3** contain signals of additional product ions, although with rather small intensities (Table III). As expected from the similar structures of the m/z 180 ions, the CA spectra are qualitatively similar. However, the intensity distribution in these CA spectra are identical only for m/z 180 ions arising from **1a-f** (including the parent compound **1a**) and are distinguished from those of **2** and **3**.




Figure 2. Potential energy diagram of a unimolecular mass spectrometric fragmentation: $I(M)$ = ionization energy; $A(a)$ = appearance energy of ion a; ΔH_R^+ = reaction enthalpy; $\Delta H_f(M^+)$, $\Delta H_f(a)$, $\Delta H_f(X)$ = heat of formation of M , a , and X ; $\Delta H_f'(a)$ = apparent heat of formation of ion a; ϵ_h^* = thermochemical activation energy of the forward reaction; ϵ_h^* = experimental activation energy of the forward reaction; ϵ^* = non-fixed excess energy of the activated complex; ϵ_f^* = thermochemical reverse activation energy; ϵ_f^* = experimental reverse activation energy; T_B = maximum kinetic energy release; $\epsilon_{product}^*$ = excess internal energy of products; q = energy partitioning quotient.

Table IV. Ionization Energy $I(M)$, Appearance Energy $A(180^+)$ Activation Energy ϵ_h^* , Apparent Heat of Formation $\Delta H_f'(180^+)$ and Corrected Heat of Formation $\Delta H_f'(180^+)_{cor}$

substituent X	$I(M)^a$	$A-(M-X)^+^a$	$\epsilon_h^+^a$	$[\Delta H_f']^b$	$[\Delta H_f']_{cor}^b$
H (1a)	8.65	9.5 (9.3)	0.8 (0.6)	237 (232)	225 (220)
<i>o</i> -F (1b)	8.66	9.5	0.7	223	220
<i>p</i> -F	8.68				
<i>o</i> -CH ₃ (1c)	8.55	9.2	0.6	241	224
<i>m</i> -CH ₃	8.48	9.7	1.2		
<i>p</i> -CH ₃	8.45	9.8	1.3		
<i>o</i> -Cl (1d)	(8.6) ^c	9.1	0.5	245	241
<i>p</i> -Cl	8.58	9.9	1.3		
<i>o</i> -Br (1e)	(8.6) ^c	9.0	0.4	260	253
<i>p</i> -Br	8.62	9.7	1.1		
<i>o</i> -I (1f)	(8.3) ^c	8.8	0.5	269	261
<i>m</i> -OCH ₃	8.27				
<i>p</i> -OCH ₃	8.15				
<i>m</i> -CF ₃	9.02				
<i>p</i> -CF ₃	8.97				
2	8.73	9.9	1.2	248	234
3	8.90	10.0	1.1	249	235

^a eV. ^b kcal mol⁻¹. ^c Estimated from Hammett plot.

This result corroborates the conclusion, drawn from the MI spectra, that **1b-f** form identical ions a, m/z 180. Furthermore, it shows that most of the stable m/z 180 ions in the mass spectrum of **1a** are also of structure a.¹⁴ These results show that the formation of m/z 180 ions from **1a-f** mainly corresponds to the reaction **1i** → **Z** → a (Scheme II) and that this intramolecular

(14) It is seen from Table III that large differences in the CA spectra of the two types of m/z 180 ions are observed for product ions m/z 151–153 formed by unimolecular decompositions. These differences persist in the CA spectra of m/z 180 ions, generated by impact with low-energy electrons (<15 eV) and stable toward unimolecular decompositions. Hence, these differences are very likely to reflect also structural differences of the m/z 180 ions and not only different internal energies.




Figure 3. Dependence of appearance energy $A(a)$ on relative dissociation energy $\Delta D(+C-X)$.

substitution reaction can be investigated without much interference by other fragmentations at least in the case of **1b-f**.

Energetics of the Fragmentation. The potential-energy diagram of a mass-spectrometric fragmentation and the energetic quantities involved are shown schematically in Figure 2.

The ionization energies ($I(M)$) of **1a-c**, **2**, and **3** and of some para- and meta-substituted derivatives of **1a** are given in Table IV together with the appearance energies $A(m/z 180)$ of $m/z 180$ ions. The $I(M)$'s of phenyl-substituted 1-phenyl-1-(2-pyridyl)-ethylenes correlate linearly with σ_{IP}^+ constants¹⁵ ($I(M) = 0.64\sigma_{IP}^+ + 8.7$). The position of the substituent at the phenyl group has only a negligible influence on the $I(M)$, even at the ortho position, and this observation has been used to estimate the $I(M)$'s of **1d-f**, which have not been determined experimentally because of the low intensities of the molecular ions.

The $A(m/z 180^+)$ values decrease in the series **1b-f** with the difference of the dissociation energies $\Delta D(+C-X)$ of the substituent in the molecular ions¹⁶ (Figure 3). The exception is the parent compound **1a**. Although $A(m/z 180^+)$ of **1a** is below that of **2** and **3**, it appears to be too large by 0.2 eV to follow the correlation. One usually observes some mean value by the experimental method used, if isobaric fragment ions with different structures and different appearance energies are formed.¹⁷ Hence the deviation of $A(m/z 180^+)$ from **1a** can be explained by the formation of ions b with a larger $A(m/z 180^+)$ besides ions a, in line with the results of the structure investigations of $m/z 180$ ions. In the following discussion data calculated with $A(a) = 9.3$ eV from **1a** are therefore included for comparison.

The apparent heat of formation of ion a, $\Delta H_f^+(a)$, can be calculated by eq 1, using the $A(a)$ from **1a-f**, the $\Delta H_f(M)$ of the neutral precursor molecules, and the $\Delta H_f(X\cdot)$ of the radicals¹⁸ lost during the fragmentation. $\Delta H_f(M)$'s have been obtained by incremental methods¹⁹ using $\Delta H_f(1,1$ -diphenylethylene) = 58.7 ± 1.1 kcal mol⁻¹²⁰ as a starting point. $\Delta H_f^+(a)$ (Table IV) includes the activation energy of the reverse reaction, ϵ_r^* , which can be (partly) corrected for by the maximum amount of kinetic energy, T_B , released during the reaction according to eq 2.

$$\Delta H_f^+(a) = A(a) + \Delta H_f(M) - \Delta H_f(X\cdot) = \Delta H_f(a) + \epsilon_r^* \quad (1)$$

$$\Delta H_f^+(a)_{cor} = \Delta H_f^+(a) - T_b \geq \Delta H_f(a) \quad (2)$$

(15) (a) Harrison, A. G.; Kebarle, P.; Lossing, F. P. *J. Am. Chem. Soc.* **1961**, *83*, 777. (b) Bursey, M. M. *Org. Mass Spectrom.* **1968**, *1*, 31. (c) Bentley, T. W.; Johnstone, R. A. W. *J. Chem. Soc. B* **1971**, 263.

(16) $D(+C-X)$ corresponds to the reaction $X\text{C}_6\text{H}_4\text{C}(\text{CH}_2)\text{C}_5\text{H}_3\text{N}^+ \cdot \rightarrow \text{C}_6\text{H}_4\text{C}(\text{CH}_2)\text{C}_5\text{H}_3\text{N} + X\cdot$. Since $\Delta H_f^+(\text{C}_6\text{H}_4\text{C}(\text{CH}_2)\text{C}_5\text{H}_3\text{N}^+)$ of the ion (with a positive charge at one of the C atoms of the phenyl group) is not known, the dissociation energies $\Delta D(+C-X)$ relative to the unsubstituted **1a**⁺ ($X = \text{H}$) have been calculated by $\Delta D(+C-X) = \Delta H_f(X\cdot) - \Delta H_f(\text{H}) + \Delta H_f(\text{1a}^+) - \Delta H_f(\text{1a}^+)$.

(17) Ocolowitz, J. L.; Cerimele, B. J.; Brown, P. *Org. Mass Spectrom.* **1974**, *8*, 61.

(18) Rosenstock, H. M.; Draxl, K.; Steiner, B. W.; Herron, J. T. *J. Phys. Chem. Ref. Data, Suppl. 1* **1977**, 6.

(19) Benson, S. W.; Cruickshank, F. R.; Golden, D. M.; Haugen, G. R.; O'Neal, H. E.; Rodgers, A. S.; Shaw, R.; Walsh, R. *Chem. Rev.* **1969**, *69*, 279.

(20) Cox, J. D.; Pilcher, G. "Thermochemistry of Organic and Organometallic Compounds"; Academic Press: New York, 1970.




Figure 4. Dependence of apparent heat of formation $\Delta H_f^+(a)$ on relative dissociation energy $\Delta D(+C-X)$.

Table V. Reaction Enthalpy ΔH_R^+ , Reverse Activation Energy ϵ_r^* , Released Kinetic Energy T , Energy Partitioning Quotient q , and Position of Transition State X_o^* of the Intramolecular Substitution of 1-Phenyl-1-(2-pyridyl)ethylene Molecular Ions

	ΔH_R^+ ^a	ϵ_r^* ^a	T_{22} ^b	T_{50} ^b	T_{22}/T_{50}	T_B ^a	q	X_o^*
1a	+2	17 (12)	420	300	1.40	12	0.71 (1.0)	0.53 (0.54)
1b	+14	3	87	42	2.07	3	1.0	0.85
1c	-6	21	540	326	1.66	17	0.81	0.42
1d	-13	25	110	50	2.20	4	0.16	0.32
1e	-30	40	192	89	2.16	7	0.18	0.19
1f	-38	49	226	105	2.15	8	0.21	0.18

^a kcal mol⁻¹. ^b meV.

Figure 4 shows a linear correlation between $\Delta H_f^+(a)$ and the relative dissociation energy $\Delta D(+C-X)$ which is expected if the same ion a, $m/z 180$, with a constant $\Delta H_f^+(a)$ arises from the fragmentations of molecular ions **1a-f**.²¹ Again $\Delta H_f^+(a)$ from **1a** fits the correlation much better if it is calculated from the corrected value of $A(m/z 180^+)$. No constant value of $\Delta H_f^+(a)$ is obtained after the correction of the $\Delta H_f^+(a)$ by T_B values, but the $\Delta H_f^+(a)_{cor}$ approach the value 220 kcal mol⁻¹, suggesting $\Delta H_f^+(a) \leq 220$ kcal mol⁻¹. This agrees very well with $\Delta H_f^+(a) = 221 \pm 5$ kcal mol⁻¹, which has been calculated independently by the thermochemical data of the hypothetical reaction sequence discussed in the Appendix. Hence $\Delta H_f^+(a) = 220$ kcal mol⁻¹ has been accepted for the further calculations.

Energy Partitioning during the Fragmentation. The kinetic energies, T , released during the formation of ion a from molecular ions **1a-f** and calculated from the peak width at 22 (T_{22}) and 50% (T_{50}) of the peak height or at the base (T_B) of the signal are given in Table V together with ϵ_r^* and the enthalpy of reaction ΔH_R^+ ; ϵ_r^* and ΔH_R^+ have been calculated according to eq 3 and 4, respectively.

$$\epsilon_r^* = A(m/z 180^+) + \Delta H_f(M) - \Delta H_f(a) - \Delta H_f(X\cdot) \quad (3)$$

$$\Delta H_R^+ = \Delta H_f(a) + \Delta H_f(X\cdot) - \Delta H_f(M)^+ \quad (4)$$

With the exception of the loss of the fluoro substituent from **1b**⁺, which is associated with only a small amount of ϵ_r^* , large values of ϵ_r^* are obtained for the loss of ortho substituents from the molecular ions of all other compounds investigated, increasing especially for the reactions of **1e**⁺ and **1f**⁺. However, the large values of ϵ_r^* are reflected in correspondingly large values of T only in case of the reactions of **1a**⁺ and **1c**⁺, but neither in the ϵ_r^* values nor in the peak shapes of the corresponding signals in the MIKE spectra of **1d**, **1e**, and **1f**. These signals are "flat-topped" in the case of **1a** and **1c**, indicating a nonstatistical distribution of ϵ_r^* during the kinetic-energy release, whereas the ratio T_{22}/T_{50} is close

(21) Schaldach, B.; Grützmaier, H. F. *Int. J. Mass Spectrom. Ion Phys.* **1979**, *31*, 271.

to the value 2.16 expected for a Gaussian peak shape²² and indicating a statistical distribution of the kinetic energies released in the case of **1d-f**. The difference in the reactions between the two classes of derivatives is most clearly seen by a comparison between **1c** and **1d**. For both compounds similar values of ϵ_r^* have been calculated (21 and 25 kcal mol⁻¹, respectively). Whereas a broad, "flat-topped" signal is observed in the MIKE spectra of **1c** for the loss of a methyl group indicating a large value of T , the Gaussian-shaped signal in the MIKE spectrum of **1d** for the loss of the chloro substituent is rather sharp, corresponding to a small kinetic-energy release.

The two classes of derivatives (and reactions) are clearly distinguished by the energy partitioning quotient q , which has been calculated with the aid of the maximum value T_B according to $q = T_B/\epsilon_r^*$. A $q = 0.71-1.0$ observed for the reactions of **1a-c** shows that all or most of ϵ_r^* is released as T during the dissociation of a H, F, and CH₃ substituent, respectively, from the molecular ions. However, q drops dramatically and abruptly to 0.16-0.21 by changing the substituent into a Cl, Br, or I atom.

A wide range of kinetic energies T and of energy partitioning quotients q has been observed for mass-spectrometric fragmentations and several theoretical models have been used to correlate the kinetic-energy release with the transition state of the reactions.^{2,4,5,6,23} One successful approach has been to calculate the potential-energy surface of the reaction and to follow the movement of the reacting system across this surface.^{5,6} However, a reliable application of these methods to reactions of large odd-electron organic ions is difficult. Hence it is of interest to look for more simple concepts which link the kinetic-energy release to certain properties of the reaction transition states.

One of these properties is seen in the "tightness" of the activated complex since a correlation appears to exist between q and the ring size of the transition state.^{6,23} However, this effect cannot be responsible for the variation of q in the reaction series studied here, because all reactions occur via five-membered transition states with probably very similar geometry. Similarly, the number of internal degrees of freedom in the reacting ions, which is also known to influence the amount of T ,^{23a-c} is not very different for **1a-f**.²⁴

A characteristic feature of the substitution reaction of **1a-f** is the slight variation of the activation energy ϵ_h^* with the dissociation energy $D(^{\circ}\text{C-X})$, while the enthalpy of reaction ΔH_R^+ changes by more than 50 kcal mol⁻¹ from endothermic or nearly thermoneutral reactions of **1a-e** to strongly exothermic reactions of **1e,f** (Table V). According to the Hammond postulate²⁵ the position of a transition state on the reaction coordinate is different for endo- and exothermic elementary reactions, respectively, with (nearly) constant activation energies. Hence, the sharp decrease in q in spite of increasing ϵ_r^* in the reaction series appears to be due to different positions of the individual transition states on the reaction coordinate.

According to Miller⁹ the position X_0^* of a transition state is determined by the height of the potential-energy barrier U^* (here ϵ_h^*) and the potential energy U_f of the reaction (here ΔH_R^+):

$$X_0^* = \frac{1}{2 - U_f/U^*} = \frac{1}{2 - \Delta H_R^+/\epsilon_h^*}$$

(22) Elder, J. F.; Beynon, J. H.; Cooks, R. G. *Org. Mass Spectrom.* **1976**, *11*, 415.

(23) (a) Beynon, J. H.; Bertrand, M.; Cooks, R. G. *J. Am. Chem. Soc.* **1973**, *95*, 1739. (b) Kim, K. C.; Beynon, J. H.; Cooks, R. G. *J. Chem. Phys.* **1974**, *61*, 1305. (c) Boyd, R. K.; Beynon, J. H. *Int. J. Mass Spectrom. Ion Phys.* **1977**, *23*, 163. (d) Elder, J. F.; Beynon, J. H.; Cooks, R. G. *Org. Mass Spectrom.* **1975**, *10*, 273. (e) Florêncio, H.; Vijfhuizen, P. C.; Heerma, W.; Dijkstra, G. *Ibid.* **1979**, *14*, 337.

(24) The different peak shapes of the signals of the corresponding metastable transitions hint to a different origin of T . The statistical distribution of T associated with a Gaussian peak shape and the small but nearly constant $q = 0.18 \pm 0.03$ for **1d-f** probably indicate that most of T stems from the ϵ^* part of ϵ_r^* (see Figure 2). In contrast, obviously ${}^0\epsilon_r^*$ is the main origin of T in the case of **1a** and **1c**, as indicated by the large values of q and the "flat-topped" form of the signals.

(25) Hammond, G. S. *J. Am. Chem. Soc.* **1955**, *77*, 334.




Figure 5. Hammond diagram of potential energy.


The positions $X_0^* = 0$ and $X_0^* = 1$ correspond to the starting and final states of the reaction, respectively. The values of X_0^* for the fragmentations of the molecular ions **1a-f** are given in Table V. They vary parallel to ΔH_R^+ from 0.85 for **1b** ("late" transition state) to 0.19 and 0.18 for **1e** and **1f** ("early" transition states). Comparing X_0^* and q in the series (Table V and Figure 5) large energy partitioning quotients of $q = 0.7-1.0$ are associated with symmetrical or late positions ($X_0^* = 0.4-1.0$) of the transition state while small energy partitioning quotients $q = 0.16-0.21$ belong to early positions ($X_0^* < 0.4$). Obviously in the first class of reactions ($X_0^* > 0.4$, late transition states) most of the energy of the transition state is accumulated in the stretched C-X bond and is released as kinetic energy of the product if the stretching of this bond proceeds toward dissociation. In the second class ($X_0^* < 0.4$, early transition states) only a minor part of the transition-state energy has to be accumulated in the dissociating C-X bond. Hence only a small part is released as kinetic energy of the dissociation products.

The present results for a correlation between energy partitioning and position of transition states fit very well into the fundamental concepts of reaction dynamics, which have been developed by Polanyi,²⁶ for the partitioning of ϵ_r^* into kinetic and vibrational energy during bimolecular reactions. From this concept it is predicted that during reactions occurring on an "attractive" potential energy surface with the energy barrier at an early position along the reaction path (reactant-like configuration of the activated complex) a vibrational excitation of the products will be favored. However, most of ${}^0\epsilon_r^*$ will be released as kinetic energy during reactions with "repulsive" energy surfaces with a late position of the transition state (product-like configuration of the activated complex). Moreover, this concept predicts that a change between the two modes of energy partitioning will take place quite suddenly, if the crest of the energy barrier is moved along the reaction path from the entry valley to the exit valley of the reaction.²⁶ This is exactly what has been observed for the series of intramolecular substitution reactions of the molecular ions **1a-f**, $X_0^* \approx 0.4$ being obviously the critical position of the transition state.

Finally, it should be mentioned that a similar behavior on energy partitioning has been observed by us recently for intramolecular substitutions of the molecular ions of 2-benzoylpyridines¹ and of benzylacetones.²¹ Although the results are somewhat disturbed by side reactions and are not as clear as for the reactions of the 1-phenyl-1-(2-pyridyl)ethylenes, a change of the q values is also observed at a critical value $X_0^* \approx 0.4$. Thus this change in the mode of energy partitioning appears to be a fundamental property of the intramolecular aromatic substitution reactions of these molecular ions, and mass-spectrometric techniques seem to be well

(26) Polanyi, J. C. *Acc. Chem. Res.* **1972**, *5*, 161.

Scheme III



sued to reveal the characteristics of such elementary reactions of complex organic ions.

Experimental Section

The mass spectra were obtained with a Varian MAT 311A mass spectrometer. Experimental conditions: accelerating voltage, 3 kV; electron energy, 70 eV; emission current, 2 mA; ion source temperature, 150 °C; ion source pressure, $< 2 \times 10^{-6}$ Torr; direct insertion of the sample and sample temperature, < 60 °C.

The kinetic energy T released during the fragmentations has been determined from the appropriate metastable transitions observed on the same instrument and similar experimental conditions by scanning the voltage across the electrostatic analyzer (second field-free region). T_{50} and T_{22} have been calculated from the peak width at the corresponding peak height.³ To determine T_B the peaks of the metastable transitions have been approximated by either a triangle or a trapezoid, and the width of the base line has been used for the calculations. All values have been corrected for the width of the main beam.²⁷ The reproducibility of the T values is $\pm 10\%$.

The ionization energies and appearance energies were determined from semilogarithmic plots of ion efficiency curves,²⁸ measured with a modified Vacuum Generators MM 12B mass spectrometer (accelerating voltage, 4 kV; electron emission current, 20 μA ; ion source temperature, 150 °C; direct insertion of the sample, repeller potential 0 V), using CH_3I ($I(M) = 9.53$ eV)¹⁸ as a reference. The values given in the tables are the mean of at least three independent measurements. The reproducibility is $< \pm 0.1$ eV.

The MI spectra and CA spectra were obtained with a Vacuum Generators ZAB-2F mass spectrometer, modified for measurements at low electron energies and equipped with collision chambers in the first and

second field-free region. He as collision gas was introduced into the appropriate collision chamber at such a rate that the intensity of the main beam dropped to 10% of its original value. With the exception of the $(\text{F}-\text{H})^+$ ions the reproducibility of the MI and CA spectra is $< \pm 10\%$.

The 1-phenyl-1-pyridylethylenes (1a-f, 2, and 3) were synthesized via the corresponding 1-methyl-1-phenyl-1-pyridylcarbinols, obtained by reaction of 2-pyridyllithium with the (substituted) benzoylpyridine by standard procedures. The carbinols were purified by vacuum distillation or recrystallization from petroleum ether (bp 60–70 °C) and dehydrated by dissolving in $\text{CH}_3\text{COOH}/\text{H}_2\text{SO}_4$ at room temperature. After the dehydration was complete (controlled by TLC), the reaction mixture was diluted with water and neutralized with 2 N NaOH. The mixture was extracted with ether. The 1-phenyl-1-pyridylethylenes were purified by column chromatography (silica, benzene/ethyl acetate, 10:1), yield 10–65%.

Acknowledgments. We are grateful to the Fonds der Chemischen Industrie for financial support of this work.

Appendix. Estimation of $\Delta H_f(a)$ by Thermochemical Data

$\Delta H_f(a)$ can be calculated from the enthalpy changes of the reaction steps a–d of the hypothetical reaction sequence of Scheme III.

$$\text{PA}(\mathbf{1a}) = \Delta H_f(\mathbf{1a}) + \Delta H_f(\text{H}^+) - \Delta H_f(\text{x}) \quad (\text{a})$$

$$D(\text{N}-\text{H}) = \Delta H_f(\text{y}) + \Delta H_f(\text{H}\cdot) - \Delta H_f(\text{x}) \quad (\text{b})$$

$$D(\text{C}_{\text{ar}}-\text{H}) = \Delta H_f(\text{z}) + \Delta H_f(\text{H}\cdot) - \Delta H_f(\text{y}) \quad (\text{c})$$

$$D(^+\text{N}-\text{Ph}) = \Delta H_f(\text{z}) - \Delta H_f(\text{a}) \quad (\text{d})$$

Equations a–d are combined to

$$\Delta H_f(a) = \Delta H_f(\mathbf{1a}) + \Delta H_f(\text{H}^+) - 2\Delta H_f(\text{H}\cdot) - \text{PA}(\mathbf{1a}) + D(\text{C}_{\text{ar}}-\text{H}) + [D(^+\text{N}-\text{H}) - D(^+\text{N}-\text{Ph})] \quad (\text{e})$$

$\Delta H_f(\mathbf{1a}) = 70$ kcal mol⁻¹ has been calculated from increments¹⁹ using $\Delta H(1,1\text{-diphenylethylene}) = 58.7 \pm 1.1$ kcal mol⁻¹²⁰ as a starting point. $D(\text{C}_{\text{ar}}-\text{H}) = 104$ kcal mol⁻¹ corresponds to the homolytic dissociation energy of a C–H bond in benzene;²⁹ $\Delta H_f(\text{H}^+) = 365.2$ ¹⁸ and $\Delta H_f(\text{H}\cdot) = 52.1$ kcal mol⁻¹¹⁸ are accurately known. From the remaining quantities of eq e the proton affinity of 1-phenyl-1-(2-pyridyl)ethylene, PA(1a), is set equal to $\text{PA}(\text{pyridine}) = 219$ kcal mol⁻¹³⁰ since the styryl substituent is not expected to alter the proton affinity due to the lone electron pair at the N atom of pyridine very much.

The homolytic dissociation energies $D(^+\text{N}-\text{H})$ and $D(^+\text{N}-\text{Ph})$ in the pyridinium ions x and a, respectively, are not known. Fortunately, only the difference between both dissociation energies has to be known, which is not large. By the use of relevant thermochemical data $[D(^+\text{N}-\text{H}) - D(^+\text{N}-\text{Ph})] = 5$ kcal mol⁻¹ has been calculated.¹ Insertion of the appropriate values in eq 3 results in $\Delta H_f(a) = 221$ kcal mol⁻¹. This is in very excellent agreement with the limiting value of 220 kcal mol⁻¹ obtained from $\Delta H_f(a)_{\text{cor}}$ by appearance-energy measurements (Table V). It should be noted that any error in $\Delta H_f(a)$ will introduce a systematic error into the calculations of ϵ_r^* and q , but will not influence the variation of q within the reaction series of 1a–f.

(27) Baldwin, M. A.; Derrick, P. J.; Morgan, R. P. *Org. Mass Spectrom.* **1976**, *11*, 440.

(28) Lossing, F. P.; Tickner, A. N.; Bryce, W. A. *J. Chem. Phys.* **1951**, *19*, 1254.

(29) Kerr, J. A. *Chem. Rev.* **1966**, *66*, 465.

(30) Freiser, B. S.; Beauchamp, J. L. *J. Am. Chem. Soc.* **1976**, *98*, 265.