

INSTITUTE OF MATHEMATICAL ECONOMICS

WORKING PAPERS


No. 324

The Dummy Paradox of the Bargaining Set

by

Bezalel Peleg and Peter Sudhölter

September 2001



University of Bielefeld

33501 Bielefeld, Germany

The Dummy Paradox of the Bargaining Set*

Bezalel Peleg[†] Peter Sudhölter[‡]

Center for Rationality and Interactive Decision Theory
The Hebrew University of Jerusalem

Abstract

By means of an example of a superadditive 0-normalized game, we show that the maximum payoff to a dummy in the bargaining set may decrease when the marginal contribution of the dummy to the grand coalition becomes positive.

We consider the weighted majority game (N, v_0) which has the tuple $(3; 1, 1, 1, 1, 1, 0)$ as a representation (see (3)). The maximum payoff to the dummy (the last player) in the bargaining set of (N, v_0) is shown to be $2/7$ (see Remark 2). If we now increase $v_0(N)$ by δ , $0 < \delta < 2/3$, then the maximum payoff to the last player in the new game, in which this player is no longer a dummy and contributes δ to N , is smaller than $2/7$ and strictly decreasing in δ (see Lemma 3).

We recall some definitions and introduce relevant notations. A (cooperative TU) *game* is a pair (N, v) such that $\emptyset \neq N$ is finite and $v : 2^N \rightarrow \mathbb{R}$, $v(\emptyset) = 0$. For any game (N, v) let

$$I(N, v) = \{x \in \mathbb{R}^N \mid x(N) = v(N) \text{ and } x^i \geq v(\{i\}) \text{ for all } i \in N\}$$

denote the set of *imputations*. (We use $x(S) = \sum_{i \in S} x^i$ for every $S \subseteq N$.) Let (N, v) be a game, $x \in I(N, v)$, and $k, l \in N$, $k \neq l$. Let

$$\mathcal{T}_{kl} = \{S \subseteq N \setminus \{l\} \mid k \in S\}.$$

An *objection* of k against l at x is a pair (P, y) satisfying

$$P \in \mathcal{T}_{kl}, \quad y(P) = v(P), \quad \text{and } y^i > x^i \text{ for all } i \in P. \quad (1)$$

We say that k can *object* against l via P , if there exists y such that (P, y) is an objection of k against l . Hence k can object against l via P , if and only if $P \in \mathcal{T}_{kl}$ and $e(P, x, v) > 0$,

*The second author was partially supported by the Edmund Landau Center for Research in Mathematical Analysis and Related Areas, sponsored by the Minerva Foundation (Germany).

[†]Email: pelegba@math.huji.ac.il

[‡]Also at the Institute of Mathematical Economics, University of Bielefeld. Email: petersud@math.huji.ac.il

where $e(S, x, v) = v(S) - x(S)$ is the excess of S at x for $S \subseteq N$. A *counter objection* to an objection (P, y) of k against l is a pair (Q, z) satisfying

$$Q \in \mathcal{T}_{lk}, z(Q) = v(Q), z^i \geq y^i \text{ for all } i \in Q \cap P \text{ and } z^j \geq x^j \text{ for all } j \in Q \setminus P. \quad (2)$$

Aumann and Maschler (1964) introduced the concepts of objections and counter objections.

An imputation $x \in I(N, v)$ is *stable* if for every objection at x there exists a counter objection. The *bargaining set* $\mathcal{M}(N, v)$ is defined by $\mathcal{M}(N, v) = \{x \in I(N, v) \mid x \text{ is stable}\}$. The bargaining set was introduced by Davis and Maschler (1967).

Player $i \in N$ is a *dummy* of (N, v) if $v(S \cup \{i\}) = v(S) + v(\{i\})$ for all $S \subseteq N \setminus \{i\}$. The game (N, v) is *superadditive* if $v(S) + v(T) \leq v(S \cup T)$ for all $S \subseteq N$ and $T \subseteq N \setminus S$.

Remark 1. Let (N, v) be a game. We recall that the core of (N, v) is the set $\mathcal{C}(N, v) = \{x \in I(N, v) \mid e(S, x, v) \leq 0 \text{ for all } S \subseteq N\}$. Also we remark (see [2]) that $\mathcal{C}(N, v) \subseteq \mathcal{M}(N, v)$.

In the sequel let $N = \{1, \dots, 6\}$ and (N, v_0) be the weighted majority game mentioned above. That is, $v_0(S)$, $S \subseteq N$, satisfies the following equation:

$$v_0(S) = \begin{cases} 0 & , \text{ if } |S \setminus \{6\}| \leq 2 \\ 1 & , \text{ if } |S \setminus \{6\}| \geq 3 \end{cases} \quad (3)$$

Then (N, v_0) is a superadditive game and player 6 is a dummy. Also, for every $\delta \in \mathbb{R}, \delta > 0$, let (N, v_δ) be the game which differs from (N, v_0) only inasmuch as $v_\delta(N) = 1 + \delta$.

If $0 \leq \delta \leq 2/3$, then define $x_\delta \in \mathbb{R}^N$ by

$$x_\delta^j = \frac{1}{7} + \frac{2}{7}\delta \text{ for } j \in N \setminus \{6\} \text{ and } x_\delta^6 = \frac{2}{7} - \frac{3}{7}\delta. \quad (4)$$

If $\delta \geq 2/3$, then define $x_\delta \in \mathbb{R}^N$ by

$$x_\delta^j = \frac{1}{3} \text{ for } j \in N \setminus \{6\} \text{ and } x_\delta^6 = \delta - \frac{2}{3}. \quad (5)$$

Remark 2. For every $\delta \geq 0$, $x_\delta \in \mathcal{M}(N, v_\delta)$.

Proof: Clearly $x_\delta \in I(N, v_\delta)$. If $\delta \geq 2/3$, then $x_\delta \in \mathcal{C}(N, v_\delta)$, thus $x_\delta \in \mathcal{M}(N, v_\delta)$ by Remark 1. Now we assume $0 \leq \delta < 2/3$. Then $\mathcal{C}(N, v_\delta) = \emptyset$. Let $k, l \in N$, $k \neq l$, and let (P, y) be an objection of k against l at x_δ . By (1), $|P \setminus \{6\}| \geq 3$. If $l \neq 6$ and $k \neq 6$, then let $Q = (P \setminus \{k\}) \cup \{l\}$. If $k = 6$, then there exists $i \in P$ and let $Q = (P \setminus \{k, i\}) \cup \{l\}$. If $l = 6$, then select $i \in P \setminus \{k\}$ satisfying $y^i \geq y^j$ for all $j \in P \setminus \{k\}$ and let $Q = N \setminus \{k, i\}$. Also, let $z \in \mathbb{R}^Q$ be given by

$$z^j = \begin{cases} y^j & , \text{ if } j \in Q \cap P \\ v(Q) - y(P \cap Q) - x_\delta(Q \setminus (P \cup \{l\})) & , \text{ if } j = l \\ x_\delta^j & , \text{ if } j \in Q \setminus (P \cup \{l\}). \end{cases} \quad (6)$$

Then (Q, z) is a counter objection to (P, y) .

q.e.d.

Lemma 3. Let $\delta \in \mathbb{R}_+$. If $x \in \mathcal{M}(N, v_\delta)$, then $x^6 \leq x_\delta^6$.

Proof: Let $x \in I(N, v_\delta)$ satisfy $x^6 > x_\delta^6$. It remains to show that $x \notin \mathcal{M}(N, v_\delta)$. Without loss of generality we may assume

$$x^1 \leq \dots \leq x^5. \quad (7)$$

In what follows we shall construct a justified objection of 1 against 6 via the coalition $P = \{1, 2, 3\}$. We distinguish two cases:

- (1) $\delta \geq 2/3$: Then 1 can object against 6 via P by (7) and the assumption that $x^6 > x_\delta^6$. Also, $\{2, 3, 4, 6\}$, $\{2, 3, 5, 6\}$, and $\{2, 4, 5, 6\}$ are the only coalitions in \mathcal{T}_{61} which might have a nonnegative excess at x . Now, player 2 is a member of all of them and $e(P, x, v_\delta) > e(Q, x, v_\delta)$ for all $Q \in \mathcal{T}_{61}$, thus there exists $y \in \mathbb{R}^P$ such that $y(P) = v(P)$, $y^i > x^i$ for all $i \in P$, and $y^2 - x^2 > e(Q, x, v_\delta)$ for all $Q \in \mathcal{T}_{61}$. We conclude that (P, y) is a justified objection of 1 against 6 at x_δ .
- (2) $0 \leq \delta < 2/3$: Again, 1 can object against 6 via P , because $x^6 > x_\delta^6$. Let $Q_{\{i\}}$, $i = 2, 3$, and $Q_{\{2,3\}}$ be the members of \mathcal{T}_{61} defined by

$$Q_{\{i\}} = \{i, 4, 5, 6\}, \quad i = 2, 3, \quad \text{and} \quad Q_{\{2,3\}} = \{2, 3, 4, 6\}.$$

Then

$$Q \in \mathcal{T}_{61}, \quad e(Q, x, v_\delta) \geq 0 \Rightarrow v_\delta(Q) = 1, \quad (8)$$

because $x \geq 0$ and $x^6 > 0$. Also, we have

$$Q \in \mathcal{T}_{61}, \quad v_\delta(Q) = 1 \Rightarrow e(Q, x, v_\delta) \leq e(Q_{Q \cap \{2,3\}}, x, v_\delta). \quad (9)$$

Indeed, every $Q \in \mathcal{T}_{61}$ satisfying $v_\delta(Q) = 1$, intersects $\{2, 3\}$, hence $Q_{Q \cap \{2,3\}}$ is defined. The inequality follows from (7). Also, $x \geq 0$, $x^6 > 0$, (7) - (9) imply that

$$e(P, x, v_\delta) > (e(Q, x, v_\delta))_+ \quad \text{for all } Q \in \mathcal{T}_{61}. \quad (10)$$

We claim that

$$e(P, x, v_\delta) > (e(Q_{\{2\}}, x, v_\delta))_+ + (e(Q_{\{3\}}, x, v_\delta))_+. \quad (11)$$

By (10) it suffices to show that

$$e(P, x, v_\delta) > e(Q_{\{2\}}, x, v_\delta) + e(Q_{\{3\}}, x, v_\delta), \quad (12)$$

which is equivalent to

$$1 - x(P) > 1 - x(Q_{\{2\}}) + 1 - x(Q_{\{3\}})$$

and, thus, to $-1 - x^1 + 2x(\{4, 5, 6\}) > 0$. By the observation that

$$-1 - x^1 + 2x(\{4, 5, 6\}) = -1 + x(N) - 2x^1 - x(\{2, 3\}) + x(\{4, 5, 6\}) \geq \delta + x^6 - 2x^1$$

it suffices to show that $\delta + x^6 - 2x^1 > 0$. By (7), $5x^1 + x^6 \leq 1 + \delta$, thus

$$\delta + x^6 - 2x^1 \geq \frac{3\delta + 7x^6 - 2}{5} > 0.$$

The last inequality is implied by the assumption that $x^6 > x_\delta^6 = 2/7 - (3/7)\delta$.
 Now the proof can be finished. By (10) and (11) there exists $t \in \mathbb{R}^P$ satisfying

$$\begin{aligned} t(P) &= e(P, x, v_\delta), \quad t(\{2, 3\}) > e(Q_{\{2,3\}}, x, v_\delta), \\ t^i &> (e(Q_{\{i\}}, x, v_\delta))_+, \quad i \in \{2, 3\}, \quad \text{and } t^1 > 0. \end{aligned} \tag{13}$$

Let $y = x^P + t$. By (13), (P, y) is a justified objection of 1 against 6 at x . q.e.d.

Remark 4. *The reactive bargaining set and the semi-reactive bargaining set, two variants of the bargaining set recently introduced by Granot and Maschler (1997) and Sudhölter and Potters (2001), do not show the dummy paradox. Indeed, in [4] it is shown, that both solutions, when restricted to superadditive games, satisfy the dummy property (that is, each member of the solution assigns $v(\{i\})$ to a dummy i).*

References

- [1] AUMANN, R. J. AND M. MASCHLER (1964): "The bargaining set for cooperative games," in M. Dresher, L. S. Shapley, and A. W. Tucker, eds., *Advances in Game Theory*, Princeton University Press, Princeton, NJ, pp. 443 - 476
- [2] DAVIS, M. AND M. MASCHLER (1967): "Existence of stable payoff configurations for cooperative games," in M. Shubik, ed., *Essays in Mathematical Economics in Honor of Oskar Morgenstern*, Princeton University Press, Princeton, NJ, pp. 39 - 52
- [3] GRANOT, D. AND M. MASCHLER (1997): "The reactive bargaining set: Structure, dynamics and extension to NTU games," *International Journal of Game Theory*, 26, pp. 75 - 95
- [4] SUDHÖLTER, P. AND J. A. M. POTTERS (2001): "The semireactive bargaining set of a cooperative game," forthcoming in the *International Journal of Game Theory*