






# Illustrating the importance of meta-analysing variances alongside means in ecology and evolution

Alfredo Sánchez-Tójar<sup>1</sup>  | Nicholas P. Moran<sup>1,2</sup>  | Rose E. O'Dea<sup>3</sup>  |  
Klaus Reinhold<sup>1</sup>  | Shinichi Nakagawa<sup>3</sup> 

<sup>1</sup>Department of Evolutionary Biology, Bielefeld University, Bielefeld, Germany

<sup>2</sup>Centre for Ocean Life DTU-Aqua, Technical University of Denmark, Lyngby, Denmark

<sup>3</sup>Evolution & Ecology Research Centre and School of Biological, Earth and Environmental Sciences, University of New South Wales, Sydney, NSW, Australia

## Correspondence

Alfredo Sánchez-Tójar, Universität Bielefeld, Abteilung Evolutionsbiologie, Morgenbreede 45, 33615 Bielefeld, Germany.  
Email: alfredo.tojar@gmail.com

## Funding information

This research was funded by the German Research Foundation (DFG) as part of the SFB TRR 212 (NC<sup>3</sup>)—Project numbers 316099922 and 396782608, and the Australian Research Discovery Grant, DP180100818. NPM was partially funded by the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 836937.

## Abstract

Meta-analysis is increasingly used in biology to both quantitatively summarize available evidence for specific questions and generate new hypotheses. Although this powerful tool has mostly been deployed to study mean effects, there is untapped potential to study effects on (trait) variance. Here, we use a recently published data set as a case study to demonstrate how meta-analysis of variance can be used to provide insights into biological processes. This data set included 704 effect sizes from 89 studies, covering 56 animal species, and was originally used to test developmental stress effects on a range of traits. We found that developmental stress not only negatively affects mean trait values, but also increases trait variance, mostly in reproduction, showcasing how meta-analysis of variance can reveal previously overlooked effects. Furthermore, we show how meta-analysis of variance can be used as a tool to help meta-analysts make informed methodological decisions, even when the primary focus is on mean effects. We provide all data and comprehensive R scripts with detailed explanations to make it easier for researchers to conduct this type of analysis. We encourage meta-analysts in all disciplines to move beyond the world of means and start unravelling secrets of the world of variance.

## KEYWORDS

coefficient of variation, early-life effects, opportunity for selection, parental effects, variability, variance ratio

## 1 | INTRODUCTION

'Our preoccupation with averages has blinded us to biological realities' (Hogben & Sim, 1953). Despite the exponential increase in the use of meta-analysis in recent years (Stewart, 2009; Gurevitch *et al.*, 2018), most meta-analyses have exclusively focused on the study of mean effects (using effect sizes such as response ratios and standardized mean differences between two groups; Nakagawa & Santos, 2012; Koricheva & Gurevitch, 2014). Meta-analysis is a

powerful statistical tool for integrating and quantitatively summarizing findings (i.e. effect sizes) from multiple studies tackling a common research question, and for generating new hypotheses. Yet, meta-analysts may be neglecting important biological realities by focusing on means alone.

In biological systems, variation from the mean is important to ecological and evolutionary processes. Phenotypic variance is important for how we understand and predict responses to selection using quantitative genetics because phenotypic variance is a key component of

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2020 The Authors. *Journal of Evolutionary Biology* published by John Wiley & Sons Ltd on behalf of European Society for Evolutionary Biology

heritability and, thus, of the breeder's equation (Arnold, 1992; Blows & Hoffmann, 2005; Walsh & Lynch, 2018). As such, phenotypic variance has been the focus of abundant research in biology, leading to the development of important evolutionary hypotheses (e.g. sex-chromosome hypothesis: James, 1973; reviewed in Reinhold & Engqvist, 2013), principles (Bateman's principles: Arnold, 1994; Bateman, 1948; reviewed in Janicke *et al.*, 2016) and entire research fields (e.g. animal personality: reviewed in Réale *et al.*, 2007, 2010).

Meta-analysis of variance components such as repeatability and heritability has been possible for a long time, and a few studies have done so (repeatability: Bell *et al.*, 2009; Holtmann *et al.*, 2017; heritability: Dochtermann *et al.*, 2019). However, only recent statistical advances in meta-analysis have made it possible to analyse differences in variance between groups (Nakagawa *et al.*, 2015), allowing us to test, for example, whether the opportunity of selection (variance in fitness) differs between groups. As a result, meta-analyses of variance are emerging (Supplementary material S1). For example, meta-analyses have shown that early-life dietary restriction not only affects mean longevity (English & Uller, 2016) but also increases variance in longevity (Senior *et al.*, 2017); that poor condition increases mean risk-taking behaviour but does not generally affect total phenotypic variance (except in specific contexts; see Moran *et al.*, 2020); and that sexual selection on males not only increases mean but also decreases variance in fitness-related traits (Cally *et al.*, 2019). Despite this, meta-analyses of variance are still rarely used.

In this study, we aim to promote the use of meta-analysis of variance in biology and other disciplines. We used a recently published meta-analytic data set of experimental studies (Eyck *et al.*, 2019) as a case study to test the prediction that developmental stress not only negatively affects mean trait values, but also increases total trait variance. Furthermore, we used meta-regression to test whether mean and variance effects differ across traits (e.g. behaviour, morphology, reproduction). Our meta-analysis of variance revealed developmental stress effects on variance, mostly on reproduction, and highlighted the importance of shifting some of our meta-analytic attention to the raw material for natural selection: variation.

## 2 | MATERIALS AND METHODS

### 2.1 | Data analysed

Experimental data on the effects of developmental stress on phenotype and fitness were obtained from Eyck *et al.* (2019). Before the analyses, we made modifications to the data set (see details in Supplementary material S2).

### 2.2 | Effect size calculation

We calculated two types of effect sizes and their associated sampling variances using the function 'escalc()' from the R package 'metafor' v.2.1-0 (Viechtbauer, 2010).

To study mean effects, we calculated the log response ratio (lnRR, Hedges *et al.*, 1999; also known as the ratio of means or ROM, Friedrich *et al.*, 2008). We chose lnRR instead of the standardized mean difference Cohen's *d* (Cohen, 1988) as used in Eyck *et al.* (2019), because (a) lnRR is less affected by heteroscedasticity (see results), and (b) lnRR can be readily interpreted as the percentage of change between the two groups. Nonetheless, for comparison with the original study, we conducted an additional meta-analysis of means based on a standardized mean difference effect size (see Supplementary Material S3).

To study variance effects, we calculated the log coefficient of variation ratio (lnCVR; Nakagawa *et al.*, 2015; Senior *et al.*, 2020). We chose lnCVR over the log variability ratio (lnVR; Nakagawa *et al.*, 2015; Senior *et al.*, 2020) because groups can simultaneously differ at both the mean and variance levels (mean–variance relationship, e.g. Taylor's Law; Cohen & Xu, 2015; Nakagawa & Schielzeth, 2012), such as that observed in our sample (see Supplementary Material S4), and lnCVR is designed to account for that. An alternative approach to account for mean–variance relationships would be to model group means and standard deviations (SD) using univariate random slope mixed-effects models or bivariate mixed-effects meta-analytic models (also called arm-based models). Compared with lnCVR, the latter approaches can have both advantages (e.g. incorporating interval scale data) and disadvantages (e.g. unknown sampling error covariances and complexity; Dias & Ades, 2016; Nakagawa *et al.*, 2015). Although we note that such approaches are possible, and have been successfully implemented elsewhere (Simons, 2015; O'Dea *et al.*, 2019), our choice of lnCVR was mostly driven by its simplicity and integration within established meta-analytic paradigms, and because lnCVR is easily comparable to lnRR.

Multiple treatment groups shared a common control group in 23 studies (25.8% of all studies) involving 252 effect sizes (35.8% of all effect sizes), leading to nonindependence among effect sizes (Lajeunesse, 2011). To deal with this nonindependence, we adjusted the sample size of the control groups to be equal to the original sample size of that control group divided by the number of times that control group was compared with a treatment group (Noble *et al.*, 2017). For the meta-analysis of means, all the revised effect sizes that we calculated were coded such that negative values indicate that developmental stress negatively affects fitness. That is, effect sizes were coded based on the expected relationship between the trait and fitness. For example, since fitness is expected to positively associate with body mass and immune response, no change in sign had to be implemented for those effect sizes. However, since fitness is expected to negatively associate with latency to reproduce and corticosterone levels, we inverted the sign of those effect sizes before the analyses (all decisions are available in data accessibility section). For the meta-analysis of variance, effect sizes were left unchanged as we expected an increase in variance across traits.

### 2.3 | Meta-analyses and meta-regressions

We ran two multilevel meta-analytic (i.e. intercept-only) models, one for each type of effect size, to test whether developmental

**TABLE 1** Results of the meta-analyses testing the effect of developmental stress on mean (lnRR) and variance (lnCVR) in phenotype and fitness. The results of the Egger's regression tests are also shown.

Effect size	k	Meta-analytic mean	$I^2_{\text{Obser.}}$ (%)	$I^2_{\text{Study}}$ (%)	$I^2_{\text{Species}}$ (%)	$I^2_{\text{Phylo}}$ (%)	$I^2_{\text{Total}}$ (%)	$Q_{\text{test}}$	Egger's test
lnRR	677	-0.12 [-0.26,0.02]	36.7 [25.1,45.2]	32.5 [18.5,43.4]	12.2 [0.0,25.5]	19.4 [0.7,38.9]	99.9 [99.9,100.0]	14993	-0.13 [-0.18,-0.06]
lnCVR	701	0.04 [-0.08,0.14]	45.9 [35.6,55.7]	21.5 [7.6,31.2]	19.3 [4.4,28.3]	4.1 [0.0,25.7]	91.7 [90.2,93.5]	1963	0.33 [0.16,0.49]

k = number of estimates;  $I^2$  = heterogeneity;  $Q_{\text{test}}$  = Cochran's Q test; NA = not applicable; Obser. = Observational or residual variance; Phylo = Phylogeny. Egger's test = intercept of an Egger's regression following Nakagawa and Santos (2012). Estimates shown correspond to modes and 95% Highest Posterior Density Intervals. N = 89 studies.

stress generally affects phenotype and fitness both at the mean (lnRR) and variance (lnCVR) levels, and two multilevel meta-regression models to test whether developmental stress effects differed across different types of traits. For meta-analytic models, we investigated unexplained variation across studies (after accounting for sampling variance) by estimating total and separate relative heterogeneity for each random effect ( $I^2$ ; Nakagawa & Santos, 2012; more in R script: 009\_results\_figures\_and\_tables.R), and absolute heterogeneity (Q) using the R package 'metafor' v.2.1-0 (Viechtbauer, 2010). For meta-regressions, we estimated the percentage of variance explained by the moderators ( $R^2_{\text{marginal}}$ ; Nakagawa & Schielzeth, 2013).

## 2.4 | Publication bias

We assessed publication bias—specifically small-study bias—by running a variant of Egger's regression that uses the meta-analytic residuals as the response variable, and the precision (i.e. the square root of the inverse of the sampling variance) as the moderator (Nakagawa & Santos, 2012). Publication bias occurs when specific effect sizes are overrepresented in the literature, and it is normally indicated by an overrepresentation of large effect sizes of small precision (small-study bias; Jennions *et al.*, 2013; Rothstein *et al.*, 2005). Additionally, we assessed potential temporal trends in effect sizes that could indicate a time-lag bias or decline effect by running a multilevel meta-regression that included year of publication as a z-transformed moderator (Nakagawa & Santos, 2012; Sánchez-Tójar *et al.*, 2018). A decline effect consists of decreasing support for a specific research hypothesis over time as evidence accumulates and is normally identified by effect sizes becoming smaller over time (Jennions & Møller, 2002; Koricheva & Kulinskaya, 2019).

## 2.5 | Random effects

All models included the following random effects: (a) observation ID, which represents the observational or residual variance that

needs to be explicitly modelled in a meta-analytic model, (b) study ID, which encompassed those estimates obtained within each specific study, (c) species ID, which encompassed those estimates obtained for each species, and (d) phylogeny, which consisted of a phylogenetic relatedness correlation matrix. To build the phylogeny, we searched for our species in the Open Tree Taxonomy (Rees & Cranston, 2017) and retrieved the phylogenetic relationships from the Open Tree of Life (Hinchliff *et al.*, 2015) using the R package 'rotl' v.3.0.5 (Michonneau *et al.*, 2016). We estimated branch lengths following Grafen (1989) as implemented in the function 'compute.brln()' of the R package 'ape' v.5.2 (Paradis & Schliep, 2019). The single polytomy encountered was dealt with via randomization using the function 'multi2di()' from the R package 'ape' v.5.2 (Paradis & Schliep, 2019).


We used the R programming language v.3.5.1 (R Core Team, 2018) throughout. All analyses were run in a Bayesian framework based on Stan programming language using the R package 'brms' v.2.9.0 (Bürkner, 2017; model parameters and convergence tests are available in the R code provided). Figures and tables were created using the R packages 'ggplot2' v.3.1.0 (Wickham, 2016) and 'gt' v.0.1.0 (Iannone *et al.*, 2019), respectively. All data and code are available from the data availability statement.

## 3 | RESULTS

Our final data set consisted of 704 effect sizes from 89 studies covering 56 animal species (Aves = 12 sp., Reptiles = 12 sp., Arthropods = 11 sp., Fishes = 9 sp., Amphibians = 6 sp., Mammals = 6 sp.; Figure S2).

### 3.1 | Meta-analysis of variance

Overall, developmental stress increased variance by around 4% on average, albeit uncertainty was high (Table 1, Figure 1). The effect of developmental stress on variance differed depending on the trait studied, with reproduction showing the largest increase in variance (ca. 21% on average) (Figure 2). However, the percentage of variance



**FIGURE 1** Developmental stress affects negatively the mean and slightly increases variance in trait values. Points and associated error bars correspond to posterior modes and 95% highest posterior density intervals (HPDI) from the meta-analyses. The posterior distributions with vertical lines indicating the median are plotted on top of their respective modes and 95% HPDI

explained by the trait moderator was less than 1% (Table 2), indicating that most heterogeneity remained unexplained.

### 3.2 | Meta-analysis of mean

Our results showed that, on average, developmental stress negatively affected mean trait values by around 13% (Table 1, Figure 1). The meta-regression showed that developmental stress negatively affected all traits, with the strongest effects being on reproduction (ca. 21% on average) and behaviour (ca. 16% on average; Table 2, Figure 2). Nonetheless, heterogeneity remained high even after including the trait moderator (Table 2). Our additional meta-analysis based on a standardized mean difference effect size led to a meta-analytic mean that was very similar to the original study (i.e. Eyck *et al.*, 2019; see Supplementary material S3). However, our phylogenetically corrected meta-analytic mean was much more uncertain (i.e. wider credible intervals overlapping zero). Additionally, relative ( $I^2$ ) and mostly absolute ( $Q$ ) heterogeneity were lower, indicating lower unexplained variation across studies in our meta-analysis (see Supplementary material S3).

## 4 | PUBLICATION BIAS

The intercepts of the Egger's regressions were clearly different from zero, suggesting the existence of publication bias (small-study bias) at both mean and variance levels (Table 1). The meta-regressions testing for temporal trends in effect sizes showed a small effect size reduction over time at both mean and variance levels, but the trends

were uncertain and the percentage of variance explained by the moderator was essentially zero (Table 2).


## 5 | DISCUSSION

Combining a recent advance in meta-analytic methodology and a case study, we demonstrate how meta-analysis of variance can shed light on important biological processes. We showed that developmental stress not only negatively affects mean trait values, but can also increase total trait variance. Our results have also revealed that developmental stress affects reproduction most strongly, both at the mean and at the variance level. Overall, we encourage meta-analysts to focus on both mean and variance effects to unearth previously overlooked effects.

### 5.1 | Case study: developmental stress effects

Developmental stress effects on phenotype and fitness have been studied often. For example, studies have investigated the effects of different developmental stressors on morphology and coloration (Tschirren *et al.*, 2009), attractiveness (Kahn *et al.*, 2012), social network position (Boogert *et al.*, 2014), telomere dynamics (Grunst *et al.*, 2019) and fitness (Arbuthnott & Whitlock, 2018). Several reviews and meta-analyses have attempted to synthesize how different developmental stressors influence phenotype and fitness. However, the majority focused on mean effects (e.g. Nakagawa *et al.*, 2012; English & Uller, 2016; Eyck *et al.*, 2019; Macartney *et al.*, 2019), with only a few recent meta-analyses exploring the effects of specific developmental stressors on variance. For example, O'Dea *et al.* (2019) showed that experimentally increasing developmental temperature leads to an ~ 8% average increase in phenotypic variance across 43 fish species, which could facilitate adaptation to novel environments by increasing the amount of rare phenotypes in the population. Dietary restriction during development has also been shown to lead to an ~ 9% average increase of variance in longevity across 14 animal species, which may affect the strength of selection on longevity (Senior *et al.*, 2017). In contrast, lower quality/quantity diets during development have been shown to lead to an ~ 8% average decrease of variance in risk-taking behaviour across animal species, suggesting that individuals may converge on high-risk behavioural phenotype under developmental diet stress (Moran *et al.*, 2020). In all, these recent meta-analyses of variance provide good evidence suggesting that developmental stress can affect phenotypic variance and the opportunity for selection.

Our results first confirm that overall, developmental stress negatively affects mean trait values, with the strongest effects on reproduction (ca. 21%) and behaviour (ca. 16%). Furthermore, our meta-analysis of variance revealed that even when multiple developmental stressors are considered together (e.g. physiological, environmental, nutritional, etc)—as in our study, developmental stress also leads to a small increase of around 4% on average in trait variance, with that effect being



**FIGURE 2** Developmental stress affects mean and variance differently across traits, with the strongest effects being on reproduction. Points and associated error bars correspond to posterior modes and 95% highest posterior density intervals (HPDI) from the meta-regressions. The posterior distributions with vertical lines indicating the median are plotted on top of their respective modes and 95% HPDI. Point size is proportional to the number of effect sizes (see Table 2)

mostly driven by an increase in variance of around 21% on average in reproduction. Thus, the increase in variance observed in experimental versus control treatments is in agreement with the recent meta-analyses above (Senior *et al.*, 2017; O'Dea *et al.*, 2019). Furthermore, our results seem in agreement with another two recent meta-analyses of variance showing that environmental stress (i.e. not only during development) measured as single-food diets and high temperature increase variance in fitness (Senior *et al.*, 2015) and reproductive success (García-Roa *et al.*, 2018), respectively. Our results on variance in reproduction, specifically, confirm previous theoretical predictions (Martin & Lenormand, 2006) and recent experimental work (Martinossi-Allibert *et al.*, 2017). Since a recent meta-analysis showed that environmental stress can increase both genetic and residual variances (i.e. not just total phenotypic variance; Rowiński & Rogell, 2017), developmental stress could have far reaching evolutionary consequences. Overall, our study shows that developmental stress may lead to increased opportunity for selection; however, these results should be interpreted carefully as most of the heterogeneity in our models remained unexplained.

## 5.2 | Promoting meta-analysis of variance

Our results show how meta-analysing variances alongside means can unearth otherwise overlooked effects and contribute to our understanding of biological processes. Indeed, all but one (93%) of the meta-analyses of variance performed in the field of ecology and evolution (see Supplementary material S1) revealed important variance effects that otherwise would have remained unknown.

Calculating InCVR for a meta-analysis of variance requires essentially the same information needed to estimate commonly used effect size statistics for comparing means such as Hedges'  $g$  (Hedges, 1981) and InRR (Hedges *et al.*, 1999). Specifically, one simply needs the mean, SD and sample size for the two groups being

compared (Nakagawa *et al.*, 2015). Since over 60% of published meta-analyses in ecology and evolution compare means (Nakagawa & Santos, 2012; Koricheva & Gurevitch, 2014), meta-analysis of variance could be applied to most meta-analytic data sets in the field, even retrospectively.

Nonetheless, there are some limitations that meta-analysts need to know when conducting a meta-analysis of variance. First, as in the case of InRR, only ratio scale data can be used to calculate InCVR, and equations to derive InCVR from other statistics such as  $F$  or  $t$  statistics are not available. Furthermore, InCVR cannot be calculated for group-level proportional data (Nakagawa *et al.*, 2015). Second, absolute error variance will generally be larger for InCVR than for mean-based effect size statistics. This large sampling variance will generally lead to lower levels of heterogeneity in InCVR compared with mean-based effect size statistics (Table 1), and overall highlights that meta-analysing variances will usually be more data-hungry than meta-analysing means. Despite these limitations, meta-analysis of variance is rather uncomplicated, making it easy for meta-analysts to shift some of their preoccupations with averages to more variance-driven hypothesis testing and development.

Meta-analysis of variance not only can reveal important biological realities, but can also help with making informed methodological decisions. By identifying whether the compared groups show unequal variances (i.e. whether there is heteroscedasticity), meta-analysis of variance can help meta-analysts choose between effect sizes that assume homoscedasticity (e.g. Cohen's  $d$ , Cohen, 1988; Hedges'  $g$ , Hedges, 1981), and those that incorporate heteroscedasticity (e.g. standardized mean difference with heteroscedasticity or SMDH, Bonett, 2008, 2009; see Supplementary material S3). This is important because not accounting for heteroscedasticity can cause parameter misestimation in meta-analysis (Bonett, 2008, 2009). Overall, we suggest that even when variance-based hypotheses are of no interest to the researcher, meta-analysis of variance can still be used



**TABLE 2** Results of the meta-regressions testing whether the effect of developmental stress on mean (lnRR) and variance (lnCVR) differs across traits. The results of the meta-regressions assessing temporal trends in effect sizes are also shown.

Estimates	Mode [95% HPDI]	N	k
<b>InRR (k = 677)</b>			
Behaviour	-0.16 [-0.30,-0.01]	33	95
Development	-0.13 [-0.29,0.01]	17	55
Metabolism and Physiology	-0.12 [-0.27,0.02]	26	118
Morphology	-0.09 [-0.22,0.06]	51	308
Reproduction	-0.18 [-0.34,-0.04]	14	62
Survival	-0.10 [-0.28,0.10]	7	39
$R^2_{\text{marginal}} (\%) =$	0.58 [0.07,1.50]	-	-
<b>InRR (k = 677; time-lag bias test)</b>			
Intercept	-0.12 [-0.25,0.01]	-	-
Year of publication	0.01 [-0.02,0.06]	-	-
$R^2_{\text{marginal}} (\%) =$	0.00 [0.00,0.51]	-	-
<b>InCVR (k = 701)</b>			
Behaviour	0.03 [-0.15,0.17]	33	100
Development	-0.01 [-0.19,0.15]	17	55
Metabolism and Physiology	0.03 [-0.11,0.20]	26	122
Morphology	0.03 [-0.11,0.14]	51	325
Reproduction	0.18 [0.01,0.36]	14	60
Survival	0.10 [-0.18,0.34]	7	39
$R^2_{\text{marginal}} (\%) =$	0.72 [0.05,2.63]	-	-
<b>InCVR (k = 701; time-lag bias test)</b>			
Intercept	0.05 [-0.07,0.16]	-	-
Year of publication	-0.03 [-0.09,0.02]	-	-
$R^2_{\text{marginal}} (\%) =$	0.01 [0.00,1.19]	-	-

N = number of studies; k = number of estimates;  $R^2_{\text{marginal}}$  = percentage of variance explained by the moderators (Nakagawa and Schielzeth 2013). Year of publication was z-transformed. Estimates shown correspond to posterior modes and 95% Highest Posterior Density Intervals (HPDI).

as a powerful methodological tool for helping to choose the most appropriate effect size statistic.

## 6 | CONCLUSION

Our analyses on the effects of developmental stress on both mean and variance in phenotype and fitness showcase how meta-analysing variances alongside means can help unravel crucial processes. Importantly, meta-analysing variances is not limited to ecology and evolution, and can also advance disciplines such as agriculture (Knapp & Heijden, 2018), social sciences (O'Dea *et al.*, 2018) and medicine (Senior *et al.*, 2016). We have also shown how meta-analysis of variance can be used as a methodological tool to make informed decisions on how to choose effect size statistics for the study of mean effects. Overall, a holistic understanding of the world requires moving beyond the world of means to incorporate the world of variance.

## ACKNOWLEDGMENTS

We are grateful to the authors of the original publication for sharing data with us and answering our questions about the data, especially Harrison J.F. Eyck and Tim S. Jessop. We are grateful to Pietro D'Amelio for enlightening discussions about data visualization, and the reviewers and the editor for their helpful comments on the manuscript.

## AUTHORS' CONTRIBUTIONS

AST contributed to conceptualization, data curation, formal analysis, investigation, methodology, project administration, provision of resources, provision of software, validation, visualization, writing the original draft of the manuscript, and review and editing of the manuscript. NPM contributed to investigation, methodology, validation, and review and editing of the manuscript. REO contributed to conceptualization, methodology, and review and editing of the manuscript. KR contributed to funding acquisition, supervision, and review and editing of the manuscript. SN contributed to conceptualization, methodology, funding acquisition, supervision, and review and editing of the manuscript.

## COMPETING INTERESTS

We declare no competing interests.

## PEER REVIEW

The peer review history for this article is available at <https://publons.com/publon/10.1111/jeb.13661>.

## DATA AVAILABILITY STATEMENT

All data and code are available at Zenodo (<http://doi.org/10.5281/zenodo.3843018>), and brms models are available as 'RData' at the Open Science Framework (<https://doi.org/10.17605/OSF.IO/YJUA8>).

## ORCID

- Alfredo Sánchez-Tójar  <https://orcid.org/0000-0002-2886-0649>  
 Nicholas P. Moran  <https://orcid.org/0000-0002-7331-0400>  
 Rose E. O'Dea  <https://orcid.org/0000-0001-8177-5075>  
 Klaus Reinhold  <https://orcid.org/0000-0002-0249-8346>  
 Shinichi Nakagawa  <https://orcid.org/0000-0002-7765-5182>

## REFERENCES

- Arbuthnott, D., & Whitlock, M. C. (2018). Environmental stress does not increase the mean strength of selection. *Journal of Evolutionary Biology*, 31, 1599–1606. <https://doi.org/10.1111/jeb.13351>
- Arnold, S. J. (1992). Constraints on phenotypic evolution. *The American Naturalist*, 140, S85–S107. <https://doi.org/10.1086/285398>
- Arnold, S. J. (1994). Bateman's principles and the measurement of sexual selection in plants and animals. *The American Naturalist*, 144, S126–S149. <https://doi.org/10.1086/285656>
- Bateman, A. J. (1948). Intra-sexual selection in *Drosophila*. *Heredity*, 2, 349–368. <https://doi.org/10.1038/hdy.1948.21>
- Bell, A. M., Hankison, S. J., & Laskowski, K. L. (2009). The repeatability of behaviour: A meta-analysis. *Animal Behaviour*, 77, 771–783. <https://doi.org/10.1016/j.anbehav.2008.12.022>
- Blows, M. W., & Hoffmann, A. A. (2005). A reassessment of genetic limits to evolutionary change. *Ecology*, 86, 1371–1384. <https://doi.org/10.1890/04-1209>
- Bonett, D. G. (2008). Confidence intervals for standardized linear contrasts of means. *Psychological Methods*, 13, 99–109. <https://doi.org/10.1037/1082-989X.13.2.99>
- Bonett, D. G. (2009). Meta-analytic interval estimation for standardized and unstandardized mean differences. *Psychological Methods*, 14, 225–238. <https://doi.org/10.1037/a0016619>
- Boogert, N. J., Farine, D. R., & Spencer, K. A. (2014). Developmental stress predicts social network position. *Biology Letters*, 10, 20140561. <https://doi.org/10.1098/rsbl.2014.0561>
- Bürkner, P.-C. (2017). brms: An R Package for Bayesian multilevel models using Stan. *Journal of Statistical Software*, 80, 1–28. <https://doi.org/10.18637/jss.v080.i01>
- Cally, J. G., Stuart-Fox, D., & Holman, L. (2019). Meta-analytic evidence that sexual selection improves population fitness. *Nature Communications*, 10, 2017. <https://doi.org/10.1038/s41467-019-10074-7>
- Cohen, J. (1988). *Statistical power analysis for the behavioral sciences*, 2nd ed. Hillsdale, NJ: L. Erlbaum Associates.
- Cohen, J. E., & Xu, M. (2015). Random sampling of skewed distributions implies Taylor's power law of fluctuation scaling. *Proceedings of the National Academy of Sciences*, 112, 7749–7754. <https://doi.org/10.1073/pnas.1503824112>
- Dias, S., & Ades, A. E. (2016). Absolute or relative effects? Arm-based synthesis of trial data. *Research Synthesis Methods*, 7, 23–28. <https://doi.org/10.1002/jrsm.1184>
- Dochtermann, N. A., Schwab, T., Anderson Berdal, M., Dalos, J., & Royauté, R. (2019). The heritability of behavior: A meta-analysis. *Journal of Heredity*, 110, 403–410. <https://doi.org/10.1093/jhered/esz023>
- English, S., & Uller, T. (2016). Does early-life diet affect longevity? A meta-analysis across experimental studies. *Biology Letters*, 12, 20160291. <https://doi.org/10.1098/rsbl.2016.0291>
- Eyck, H. J. F., Buchanan, K. L., Crino, O. L., & Jessop, T. S. (2019). Effects of developmental stress on animal phenotype and performance: A quantitative review. *Biological Reviews*, 94, 1143–1160. <https://doi.org/10.1111/brv.12496>
- Friedrich, J. O., Adhikari, N. K., & Beyene, J. (2008). The ratio of means method as an alternative to mean differences for analyzing continuous outcome variables in meta-analysis: A simulation study. *BMC Medical Research Methodology*, 8, 32. <https://doi.org/10.1186/1471-2288-8-32>
- García-Roa, R., García-Gonzalez, F., Noble, D.W.A., & Carazo, P. (2018). Temperature as a modulator of sexual selection. *EcoEvoRxiv*. <https://doi.org/10.32942/osf.io/hqvmd>
- Grafen, A. (1989). The phylogenetic regression. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 326, 119–157. <https://doi.org/10.1098/rstb.1989.0106>
- Grunst, A. S., Grunst, M. L., Gonser, R. A., & Tuttle, E. M. (2019). Developmental stress and telomere dynamics in a genetically polymorphic species. *Journal of Evolutionary Biology*, 32, 134–143. <https://doi.org/10.1111/jeb.13400>
- Gurevitch, J., Koricheva, J., Nakagawa, S., & Stewart, G. (2018). Meta-analysis and the science of research synthesis. *Nature*, 555, 175–182. <https://doi.org/10.1038/nature25753>
- Hedges, L. V. (1981). Distribution theory for glass's estimator of effect size and related estimators. *Journal of Educational Statistics*, 6, 107–128. <https://doi.org/10.3102/10769986006002107>
- Hedges, L.V., Gurevitch, J., & Curtis, P.S. (1999). The meta-analysis of response ratios in experimental ecology. *Ecology*, 80, 1150–1156. [https://doi.org/10.1890/0012-9658\(1999\)080\[1150:TMAOR\]2.0.CO;2](https://doi.org/10.1890/0012-9658(1999)080[1150:TMAOR]2.0.CO;2)
- Hinchliff, C. E., Smith, S. A., Allman, J. F., Burleigh, J. G., Chaudhary, R., Coghill, L. M. ... Cranston, K. A. (2015). Synthesis of phylogeny and taxonomy into a comprehensive tree of life. *Proceedings of the National Academy of Sciences*, 112, 12764–12769. <https://doi.org/10.1073/pnas.1423041112>
- Hogben, L., & Sim, M. (1953). The self-controlled and self-recorded clinical trial for low-grade morbidity. *Journal of Epidemiology & Community Health*, 7, 163–179. <https://doi.org/10.1136/jech.7.4.163>
- Holtmann, B., Lagisz, M., & Nakagawa, S. (2017). Metabolic rates, and not hormone levels, are a likely mediator of between-individual differences in behaviour: A meta-analysis. *Functional Ecology*, 31, 685–696. <https://doi.org/10.1111/1365-2435.12779>
- Iannone, R., Cheng, J., & Schloerke, B. (2019). gt: Easily create presentation-ready display tables.
- James, J. W. (1973). 353. Note: Covariances between relatives due to sex-linked genes. *Biometrics*, 29, 584–588. <https://doi.org/10.2307/2529178>
- Janicke, T., Häderer, I. K., Lajeunesse, M. J., & Anthes, N. (2016). Darwinian sex roles confirmed across the animal kingdom. *Science Advances*, 2, e1500983. <https://doi.org/10.1126/sciadv.1500983>
- Jennions, M. D., & Møller, A. P. (2002). Relationships fade with time: A meta-analysis of temporal trends in publication in ecology and evolution. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, 269, 43–48. <https://doi.org/10.1098/rspb.2001.1832>
- Jennions, M., Lortie, C.J., Rosenberg, M.S., & Rothstein, H. (2013). Publication and related biases. *Handbook of meta-analysis in ecology & evolution* (pp. 207–236). Princeton, NJ: Princeton University Press.
- Kahn, A. T., Livingston, J. D., & Jennions, M. D. (2012). Do females preferentially associate with males given a better start in life? *Biology Letters*, 8, 362–364. <https://doi.org/10.1098/rsbl.2011.1106>
- Knapp, S., & van der Heijden, M.G.A. (2018). A global meta-analysis of yield stability in organic and conservation agriculture. *Nature Communications*, 9, 1–9. <https://doi.org/10.1038/s41467-018-05956-1>
- Koricheva, J., & Gurevitch, J. (2014). Uses and misuses of meta-analysis in plant ecology. *Journal of Ecology*, 102, 828–844. <https://doi.org/10.1111/1365-2745.12224>
- Koricheva, J., & Kulinskaya, E. (2019). Temporal instability of evidence base: A threat to policy making? *Trends in Ecology & Evolution*, 34, 895–902. <https://doi.org/10.1016/j.tree.2019.05.006>
- Lajeunesse, M. J. (2011). On the meta-analysis of response ratios for studies with correlated and multi-group designs. *Ecology*, 92, 2049–2055. <https://doi.org/10.1890/11-0423.1>

- Macartney, E. L., Crean, A. J., Nakagawa, S., & Bonduriansky, R. (2019). Effects of nutrient limitation on sperm and seminal fluid: A systematic review and meta-analysis. *Biological Reviews*, *94*, 1722–1739. <https://doi.org/10.1111/brv.12524>
- Martin, G., & Lenormand, T. (2006). A general multivariate extension of Fisher's geometrical model and the distribution of mutation fitness effects across species. *Evolution*, *60*, 893–907. <https://doi.org/10.1111/j.0014-3820.2006.tb01169.x>
- Martinossi-Allibert, I., Arnqvist, G., & Berger, D. (2017). Sex-specific selection under environmental stress in seed beetles. *Journal of Evolutionary Biology*, *30*, 161–173. <https://doi.org/10.1111/jeb.12996>
- Michonneau, F., Brown, J.W., & Winter, D.J. (2016). rotI: An R package to interact with the Open Tree of Life data. *Methods in Ecology and Evolution*, *7*, 1476–1481. <https://doi.org/10.1111/2041-210X.12593>
- Moran, N.P., Sánchez-Tójar, A., Schielzeth, H., & Reinhold, K. (2020). Poor nutritional condition promotes high-risk behaviours: A systematic review and meta-analysis. <https://doi.org/10.32942/osf.io/xsehd>
- Nakagawa, S., & Santos, E.S.A. (2012). Methodological issues and advances in biological meta-analysis. *Evolutionary Ecology*, *26*, 1253–1274. <https://doi.org/10.1007/s10682-012-9555-5>
- Nakagawa, S., & Schielzeth, H. (2012). The mean strikes back: Mean-variance relationships and heteroscedasticity. *Trends in Ecology & Evolution*, *27*, 474–475. <https://doi.org/10.1016/j.tree.2012.04.003>
- Nakagawa, S., & Schielzeth, H. (2013). A general and simple method for obtaining R<sup>2</sup> from generalized linear mixed-effects models. *Methods in Ecology and Evolution*, *4*, 133–142. [https://doi.org/10.1111/j.2041-210x.2012.00261.x@10.1111/\(ISSN\)2041-210X.STATSTOO](https://doi.org/10.1111/j.2041-210x.2012.00261.x@10.1111/(ISSN)2041-210X.STATSTOO)
- Nakagawa, S., Lagisz, M., Hector, K. L., & Spencer, H. G. (2012). Comparative and meta-analytic insights into life extension via dietary restriction. *Aging Cell*, *11*, 401–409. <https://doi.org/10.1111/j.1474-9726.2012.00798.x>
- Nakagawa, S., Poulin, R., Mengersen, K., Reinhold, K., Engqvist, L., Lagisz, M., ... Senior, A. M. (2015). Meta-analysis of variation: Ecological and evolutionary applications and beyond. *Methods in Ecology and Evolution*, *6*, 143–152. <https://doi.org/10.1111/2041-210X.12309>
- Noble, D. W. A., Lagisz, M., O'dea, R. E., & Nakagawa, S. (2017). Nonindependence and sensitivity analyses in ecological and evolutionary meta-analyses. *Molecular Ecology*, *26*, 2410–2425. <https://doi.org/10.1111/mec.14031>
- O'Dea, R. E., Lagisz, M., Jennions, M. D., & Nakagawa, S. (2018). Gender differences in individual variation in academic grades fail to fit expected patterns for STEM. *Nature Communications*, *9*, 1–8. <https://doi.org/10.1038/s41467-018-06292-0>
- O'Dea, R. E., Lagisz, M., Hendry, A. P., & Nakagawa, S. (2019). Developmental temperature affects phenotypic means and variability: A meta-analysis of fish data. *Fish and Fisheries*, *20*, 1005–1022. <https://doi.org/10.1111/faf.12394>
- Paradis, E., & Schliep, K. (2019). ape 5.0: An environment for modern phylogenetics and evolutionary analyses in R. *Bioinformatics*, *35*, 526–528. <https://doi.org/10.1093/bioinformatics/bty633>
- R Core Team. (2018). *R: A language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing.
- Réale, D., Reader, S. M., Sol, D., McDougall, P. T., & Dingemans, N. J. (2007). Integrating animal temperament within ecology and evolution. *Biological Reviews*, *82*, 291–318. <https://doi.org/10.1111/j.1469-185X.2007.00010.x>
- Réale, D., Dingemans, N. J., Kazem, A. J. N., & Wright, J. (2010). Evolutionary and ecological approaches to the study of personality. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *365*, 3937–3946. <https://doi.org/10.1098/rstb.2010.0222>
- Rees, J., & Cranston, K. (2017). Automated assembly of a reference taxonomy for phylogenetic data synthesis. *Biodiversity Data Journal*, *5*, e12581. <https://doi.org/10.3897/BDJ.5.e12581>
- Reinhold, K., & Engqvist, L. (2013). The variability is in the sex chromosomes. *Evolution*, *67*, 3662–3668. <https://doi.org/10.1111/evo.12224>
- Rothstein, H. R., Sutton, A. J., & Borenstein, M. (2005). Publication bias in meta-analysis: Prevention, assessment and adjustments. John Wiley & Sons, Ltd. <https://doi.org/10.1002/0470870168.ch1>
- Rowiński, P. K., & Rogell, B. (2017). Environmental stress correlates with increases in both genetic and residual variances: A meta-analysis of animal studies. *Evolution*, *71*, 1339–1351. <https://doi.org/10.1111/evo.13201>
- Sánchez-Tójar, A., Nakagawa, S., Sánchez-Fortún, M., Martin, D.A., Ramani, S., Girndt, A., ... Schroeder, J. (2018). Meta-analysis challenges a textbook example of status signalling and demonstrates publication bias. *Elife*, *7*, e37385. <https://doi.org/10.7554/eLife.37385>
- Senior, A. M., Nakagawa, S., Lihoreau, M., Simpson, S. J., & Raubenheimer, D. (2015). An overlooked consequence of dietary mixing: A varied diet reduces interindividual variance in fitness. *The American Naturalist*, *186*, 649–659. <https://doi.org/10.1086/683182>
- Senior, A. M., Gosby, A. K., Lu, J., Simpson, S. J., & Raubenheimer, D. (2016). Meta-analysis of variance: An illustration comparing the effects of two dietary interventions on variability in weight. *Evolution, Medicine, and Public Health*, *2016*, 244–255. <https://doi.org/10.1093/emph/eow020>
- Senior, A. M., Nakagawa, S., Raubenheimer, D., Simpson, S. J., & Noble, D. W. A. (2017). Dietary restriction increases variability in longevity. *Biology Letters*, *13*, 20170057. <https://doi.org/10.1098/rsbl.2017.0057>
- Senior, A. M., Viechtbauer, W., & Nakagawa, S. (2020). Revisiting and expanding meta-analysis of variation: the log coefficient of variation ratio, lnCVR. *Research Synthesis Methods*. <https://doi.org/10.1002/jrsm.1423>
- Simons, M. J. P. (2015). Questioning causal involvement of telomeres in aging. *Ageing Research Reviews*, *24*, 191–196. <https://doi.org/10.1016/j.arr.2015.08.002>
- Stewart, G. (2009). Meta-analysis in applied ecology. *Biology Letters*, *6*, 78–81. <https://doi.org/10.1098/rsbl.2009.0546>
- Tschirren, B., Rutstein, A. N., Postma, E., Mariette, M., & Griffith, S. C. (2009). Short- and long-term consequences of early developmental conditions: A case study on wild and domesticated zebra finches: Life-history strategies of wild and domestic birds. *Journal of Evolutionary Biology*, *22*, 387–395. <https://doi.org/10.1111/j.1420-9101.2008.01656.x>
- Viechtbauer, W. (2010). Conducting meta-analyses in R with the metafor Package. *Journal of Statistical Software*, *36*, 1–48. <https://doi.org/10.18637/jss.v036.i03>
- Walsh, B., & Lynch, M. (2018). *Evolution and selection of quantitative traits*. New York, NY: Oxford University Press.
- Wickham, H. (2016). *ggplot2: Elegant graphics for data analysis*, 2nd ed. Use R! Springer International Publishing.

## SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section.

**How to cite this article:** Sánchez-Tójar A, Moran NP, O'Dea RE, Reinhold K, Nakagawa S. Illustrating the importance of meta-analysing variances alongside means in ecology and evolution. *J Evol Biol*. 2020;33:1216–1223. <https://doi.org/10.1111/jeb.13661>