

The icosahedral group and the homotopy of the stable mapping class group

Johannes Ebert

(Communicated by Wolfgang Lück)

Abstract. We construct an explicit generator for $\pi_3(B\Gamma_\infty^+)$, the third homotopy group of the classifying space of stable mapping class group after Quillen’s plus construction. The construction relies on a very classical action of the binary icosahedral group on some hyperelliptic Riemann surface.

1. INTRODUCTION AND RESULTS

Let F_g be an oriented closed connected surface of genus g , let $\text{Diff}^+(F_g)$ be the topological group of orientation-preserving diffeomorphisms of F_g and let $\Gamma_g := \pi_0(\text{Diff}^+(F_g))$ be the *mapping class group*. If $g \geq 2$, then all components of $\text{Diff}^+(F_g)$ are contractible [3] and so $B\Gamma_g \simeq B\text{Diff}^+(F_g)$, in other words, $B\Gamma_g$ is the base space of the universal surface bundle $E_g \rightarrow B\Gamma_g$ of genus g .

It is known that Γ_g is a perfect group [17] if $g \geq 3$ and so Quillen’s plus construction—a familiar construction from algebraic K-theory, cp. [18, p. 266]—can be applied to $B\Gamma_g$. The result is a simply-connected space $B\Gamma_g^+$ and a homology equivalence $B\Gamma_g \rightarrow B\Gamma_g^+$. By definition, $B\Gamma_g$ is aspherical (i.e. all homotopy groups but the fundamental group are trivial), but $B\Gamma_g^+$ has many nontrivial higher homotopy groups.

How can one construct elements in $\pi_n(B\Gamma_g^+)$? Suppose that M is an n -dimensional homology sphere and suppose that $\pi_1(M)$ acts on F_g by orientation preserving diffeomorphisms. Such an action yields an homomorphism $\rho : \pi_1(M) \rightarrow \Gamma_g$. Let $\lambda : M \rightarrow B\pi_1(M)$ be a classifying map for the universal covering of M . Consider the composition $B\rho \circ \lambda : M \rightarrow B\Gamma_g$. Quillen’s plus construction gives a map $M^+ \rightarrow B\Gamma_g^+$. On the other hand, any degree one map $M \rightarrow \mathbb{S}^n$ is a homology equivalence and so it induces a homotopy

Supported by a fellowship within the Postdoc-Programme of the German Academic Exchange Service (DAAD).

equivalence $M^+ \simeq \mathbb{S}^n$. We end up with a map $\mathbb{S}^n \rightarrow B\Gamma_g^+$, giving an element $\theta_{M;\rho} \in \pi_n(B\Gamma_g^+)$ which only depends on M and ρ .

The purpose of this note is to study an example of such an action when $M = \mathbb{S}^3/\hat{G}$ is the Poincaré sphere (the quotient of \mathbb{S}^3 by the binary icosahedral group \hat{G}) and $g = 14$. Here is our main result.

Theorem 1. *There exists an action of the binary icosahedral group $\hat{G} = \pi_1(M)$ on F_{14} , such that $\theta_{M;\rho}$ is a generator of $\pi_3(B\Gamma_{14}^+) \cong \mathbb{Z}/24$.*

Before we delve into the details, we explain our motivation for this apparently very specialized result. It comes from the recent homotopy theory of the *stable* mapping class group. To define the stable mapping class groups, we need to consider the mapping class groups of surfaces with boundary components. Let F be an oriented surface of genus g with b boundary components, let $\text{Diff}(F, \partial F)$ be the topological group of orientation preserving diffeomorphisms which restrict to the identity on the boundary and put $\Gamma_{g,b} := \pi_0 \text{Diff}(F; \partial)$. These groups are related by stabilization maps $\Gamma_{g,b} \rightarrow \Gamma_{g,b+1}$, $\Gamma_{g,b} \rightarrow \Gamma_{g,b-1}$, $\Gamma_{g,b} \rightarrow \Gamma_{g+1,b}$ for $b > 0$, which are obtained by gluing in pairs of pants, a disc or a torus in an appropriate way. The homological stability theorems of Harer [6], as improved by Ivanov [9], [10], assert that these maps induce isomorphisms on group homology $H_k(B\Gamma_{g,b}; \mathbb{Z})$ as long as $g \geq 2k + 2$. Let $\Gamma_{\infty,b} := \text{colim}(\dots \rightarrow \Gamma_{g,b+1} \rightarrow \Gamma_{g+1,b+1} \rightarrow \dots)$. Abbreviate $\Gamma_{\infty} := \Gamma_{\infty,0}$. Quillen's plus construction gives maps $B\Gamma_{g,1}^+ \rightarrow B\Gamma_{\infty}^+$ and $B\Gamma_{g,1}^+ \rightarrow B\Gamma_g^+$. These maps are highly connected because of Harer stability. In particular, we get an isomorphism $\phi : \pi_n(B\Gamma_g^+) \rightarrow \pi_n(B\Gamma_{\infty}^+)$ for all pairs (g, n) with $g \geq 2n + 2$, for example for $(g, n) = (14, 3)$. So Theorem 1 is really about $\pi_3(B\Gamma_{\infty}^+)$. Note that, however, the isomorphism ϕ is *not* induced from a group homomorphism $\Gamma_g \rightarrow \Gamma_{\infty}$. There is a map $B\Gamma_g \rightarrow B\Gamma_{\infty}^+$ which induces ϕ , but its construction requires the Madsen-Weiss theorem [13].

Due to the work of Tillmann, Madsen, Weiss and Galatius [20], [12], [13], [5], the homology of $B\Gamma_{\infty}$ is completely understood, at least in theory and one also gets some information about the homotopy groups of $B\Gamma_{\infty}^+$. Let us describe this story briefly.

Madsen and Tillmann [12] constructed a spectrum $\mathbf{MTSO}(2)$ and maps $\alpha_{g,b} : B\Gamma_{g,b} \rightarrow \Omega_0^{\infty} \mathbf{MTSO}(2)$ ([12] uses the notation $\mathbb{C}\mathbb{P}_{-1}^{\infty}$ for $\mathbf{MTSO}(2)$). The maps $\alpha_{g,b}$ are compatible with the gluing maps and so they define a limit map $\alpha_{\infty} : B\Gamma_{\infty} \rightarrow \Omega_0^{\infty} \mathbf{MTSO}(2)$. The rational homology of $\Omega_0^{\infty} \mathbf{MTSO}(2)$ is easy to determine (it is isomorphic to the rational homology of BU , see [12]). The very complicated homology with \mathbb{F}_p -coefficients for all primes p was computed by Galatius [5].

Madsen and Weiss showed that α_{∞} is a homology equivalence. Therefore, the plus construction $(\alpha_{\infty})^+ : B\Gamma_{\infty}^+ \rightarrow \Omega_0^{\infty} \mathbf{MTSO}(2)$ is a homotopy equivalence. Then $\beta_g = (\alpha_{\infty}^+)^{-1} \circ \alpha_g : B\Gamma_g \rightarrow \Omega_0^{\infty} \mathbf{MTSO}(2) \rightarrow B\Gamma_{\infty}^+$ induces the isomorphism ϕ above after plus construction.

The first few homotopy groups of $\Omega_0^{\infty} \mathbf{MTSO}(2)$ and hence of $B\Gamma_{\infty}^+$ are given by the following theorem.

Theorem 2.

$$\pi_k(B\Gamma_\infty^+) \cong \pi_k(\Omega_0^\infty \mathbf{MTSO}(2)) \begin{cases} 0 & \text{if } k = 1 \\ \mathbb{Z} & \text{if } k = 2 \\ \mathbb{Z}/24 & \text{if } k = 3. \end{cases}$$

This is proved in [12, p. 537], but the proof of the third isomorphism is only sketched. The proof can easily be completed using the computations of the first few stable homotopy groups of $\mathbb{C}P^\infty$ from [16, p. 199].

We should remark that the idea of defining elements of $\pi_n(X^+)$ using homology spheres is not new. For $n = 3$ and $X = B\mathrm{Gl}_\infty(\mathbb{Z})$, Jones and Westbury [11] give detailed computations. Their results could be used to prove Theorem 1, but our computations are more elementary and do not depend on [11].

Outline of the paper: In Section 2, we construct a homomorphism $\eta : \pi_3(B\Gamma_\infty^+) \rightarrow \mathbb{Q}/\mathbb{Z}$ which will be used later on to detect homotopy classes. It is closely related to the classical e -invariant $K_3(\mathbb{Z}) \rightarrow \mathbb{C}/\mathbb{Z}$ in algebraic K-theory, as we will explain in Section 3.

In Section 4, we recall some facts about the icosahedral groups and we construct the action of \hat{G} on F_{14} of Theorem 1. The computations leading to Theorem 1 are given in Section 5.

Acknowledgements: This paper is an improved version of a chapter of the author’s PhD thesis [4]. The author wants to express his thanks to his PhD advisor, Carl-Friedrich Bödigheimer, for his constant support and patience. Also, I want to thank the Max-Planck-Institute for Mathematics in Bonn for financial support during my time as a PhD student. The final writing of this paper was done while the author stayed at the Mathematical Institute of the Oxford University, which was made possible by a grant from the Deutscher Akademischer Austauschdienst.

2. AN INVARIANT $\eta : \pi_3(B\Gamma_\infty^+) \rightarrow \mathbb{Q}/\mathbb{Z}$

Our procedure to detect elements in $\pi_3(B\Gamma_\infty^+)$ is based on two theorems by Harer. The first one [8] is the isomorphism (for $g \geq 8$)

$$(3) \quad H^3(B\Gamma_g; \mathbb{Q}) = 0,$$

which can also be derived from Theorem 2. The second one is that $H^2(B\Gamma_\infty; \mathbb{Z}) \cong \mathbb{Z}$, [7]. We need a precise description of the second cohomology group.

The mapping class group Γ_g acts on the cohomology of the surface F_g preserving the intersection form. The action defines a group homomorphism $\Gamma_g \rightarrow \mathrm{Sp}_{2g}(\mathbb{Z}) \rightarrow \mathrm{Sp}_{2g}(\mathbb{R})$. Because $U(g) \subset \mathrm{Sp}_{2g}(\mathbb{R})$ is a maximal compact subgroup, $B\mathrm{Sp}_{2g}(\mathbb{R}) \simeq BU(g)$. We obtain a map $\Phi_g : B\Gamma_g \rightarrow BU(g)$.

This map classifies a complex vector bundle $V_1 \rightarrow B\Gamma_g$, which has the following alternative description. Consider the universal surface bundle $\pi : E_g \rightarrow B\Gamma_g$. One can choose a complex structure on the vertical tangent bundle $T_v E_g$ of this bundle, turning $\pi : E_g \rightarrow B\Gamma_g$ into a family of Riemann surfaces. Let V_1 be the vector bundle whose fiber over $x \in B\Gamma_g$ is the dual

of the g -dimensional vector space of holomorphic 1-forms on $\pi^{-1}(x)$. This is indeed a vector bundle once the topology is appropriately chosen, see [2]. The isomorphism class of V_1 does not depend on the choice of the complex structure because that complex structure is unique up to homotopy. The Hodge decomposition theorem shows that Φ_g is a classifying map for V_1 .

There exists a universal map $\Phi : \Omega^\infty \mathbf{MTSO}(2) \rightarrow BU$, such that $\iota \circ \Phi_g \sim \Phi \circ \alpha_g$ ($\iota : BU(g) \rightarrow BU$ is the inclusion). This is shown in [12, p. 539 f] (they denote Φ by η'). Details can also be found in [4, Ch. 6].

The map Φ can be used to obtain cohomology classes of $\Omega^\infty \mathbf{MTSO}(2)$ and hence, via α_g , on each $B\Gamma_g$. We put $\zeta_i := \alpha_g^* \Phi^* c_i$ and $\gamma_i := \alpha_g^* \Phi^* s_i$; $s_i = i! \text{ch}_i \in H^{2i}(BU; \mathbb{Z})$ the i th component of the integral Chern character. The following is the main result of [7]¹, which also can be derived from Theorem 2.

Theorem 4. *The group $H^2(B\Gamma_g; \mathbb{Z})$ is infinite cyclic, generated by ζ_1 .*

We remark that ζ_1 is the first Chern class of the determinant bundle $\det(V_1)$ of the bundle V_1 . Another result we will need was proved by Morita [15, p. 555], as a consequence of Chern-Weil theory.

Lemma 5. *The class $\gamma_{2i} \in H^{2i}(B\Gamma_\infty^+)$ is a torsion class.*

Now we want to construct a homomorphism

$$(6) \quad \eta : \pi_3(B\Gamma_\infty^+) \rightarrow \mathbb{Q}/\mathbb{Z}.$$

For any space X , let $q_n^X : \tau_{\geq n} X \rightarrow X$ be the $(n - 1)$ -connected cover of X . The space $\tau_{\geq n} X$ is $(n - 1)$ -connected and q_n^X induces an isomorphism $\pi_k(\tau_{\geq n} X) \rightarrow \pi_k(X)$ for all $k \geq n$. If X is understood, we write $q_n := q_n^X$. The assignment $X \mapsto \tau_{\geq n} X$ is a functor on the homotopy category of pointed spaces and the maps q_n^X define a natural transformation of functors $q_n : \tau_{\geq n} \rightarrow \text{id}$.

If $X = B\Gamma_\infty^+$ and $n = 3$, then $\tau_{\geq 3} B\Gamma_\infty^+$ is the homotopy fiber of $\zeta_1 : B\Gamma_\infty^+ \rightarrow K(\mathbb{Z}; 2)$ by Theorem 2 and Theorem 4 and we can view $\tau_{\geq 3} B\Gamma_\infty^+$ as the total space of the circle bundle associated with $\det(V_1)$.

Lemma 7. *The cohomology class $q_3^* \zeta_2 \in H^4(\tau_{\geq 3} B\Gamma_\infty^+; \mathbb{Z})$ is a torsion class.*

Proof. Recall the relation

$$(8) \quad s_2 = c_1^2 - 2c_2 \in H^4(BU; \mathbb{Z}).$$

Applying Φ^* yields $\gamma_2 = \zeta_1^2 - 2\zeta_2 \in H^4(B\Gamma_\infty^+; \mathbb{Z})$. Consequently, $q_3^*(\gamma_2) = q_3^*(\zeta_1^2) - 2q_3^*(\zeta_2) = 0 - 2q_3^*(\zeta_2)$. By Lemma 5, γ_2 is a torsion class. \square

Because of Theorem 2, $H^3(\tau_{\geq 3} B\Gamma_\infty^+; \mathbb{Q}) = 0$. Therefore we get an exact sequence

$$(9) \quad 0 \longrightarrow H^3(\tau_{\geq 3} B\Gamma_\infty^+; \mathbb{Q}/\mathbb{Z}) \xrightarrow{\delta} H^4(\tau_{\geq 3} B\Gamma_\infty^+; \mathbb{Z}) \longrightarrow H^4(\tau_{\geq 3} B\Gamma_\infty^+; \mathbb{Q})$$

¹This is stated falsely in [7], but Harer corrected the mistake in [6].

from the Bockstein exact sequence. Thus there exists a unique

$$\bar{\zeta}_2 \in H^3(\tau_{\geq 3}B\Gamma_\infty^+; \mathbb{Q}/\mathbb{Z}) \text{ such that } \delta(\bar{\zeta}_2) = q_3^* \zeta_2.$$

Let $x : \mathbb{S}^3 \rightarrow B\Gamma_\infty^+$ be a map representing an element $x \in \pi_3(B\Gamma_\infty^+)$. There is a unique lift $\tilde{x} : \mathbb{S}^3 \rightarrow \tau_{\geq 3}B\Gamma_\infty^+$ of x . We define a homomorphism $\eta : \pi_3(B\Gamma_\infty^+) \rightarrow \mathbb{Q}/\mathbb{Z}$ by

$$(10) \quad \eta(x) := \langle \tilde{x}^* \bar{\zeta}_2; [\mathbb{S}^3] \rangle \in \mathbb{Q}/\mathbb{Z}.$$

3. COMPARISON WITH THE e -INVARIANT

Let us briefly describe the Adams-Quillen e -invariant, following [19]; it is a homomorphism $K_{2n-1}(\mathbb{C}_d) \rightarrow \mathbb{C}/\mathbb{Z}$. Here \mathbb{C}_d denotes the complex numbers with the discrete topology and the K -theory group is $K_n(\mathbb{C}_d) = \pi_n(BG\Gamma_\infty(\mathbb{C}_d)^+)$. Let FC be the homotopy fiber of the Chern character $ch : BU \rightarrow \prod_{n \geq 1} K(\mathbb{C}, 2n)$. By Bott periodicity, $\pi_{2k}(FC) = 0$ and $\pi_{2k+1}(FC) \cong \mathbb{C}/\mathbb{Z}$. The ‘‘identity’’ $\mathbb{C}_d \rightarrow \mathbb{C}$ is continuous and induces a map $T : B\Gamma_\infty(\mathbb{C}_d)^+ \rightarrow B\Gamma(\mathbb{C}) \simeq BU$. By Chern-Weil theory, the induced map in complex cohomology $T^* : H^*(BU; \mathbb{C}) \rightarrow H^*(B\Gamma_\infty(\mathbb{C}_d)^+; \mathbb{C})$ is trivial in positive degrees, cp. [14, Lem. 12] (the argument in *loc.cit.* generalizes easily to reductive connected groups as $GL_n(\mathbb{C})$ and the passage to the limit $n \rightarrow \infty$ is also straightforward). It follows that there exists a lift in the diagram

$$(11) \quad \begin{array}{ccc} & & FC \\ & \nearrow \tilde{T} & \downarrow \\ B\Gamma_\infty(\mathbb{C}_d)^+ & \xrightarrow{T} & BU \end{array}$$

This lift is not unique, but Chern-Weil theory defines a specific choice of a lift, cp. [11, Sec. 2]. On homotopy groups, \tilde{T} induces $e : K_{2n-1}(\mathbb{C}_d) \rightarrow \pi_{2n-1}(FC) \cong \mathbb{C}/\mathbb{Z}$, which is the Adams-Quillen e -invariant.

There is the following alternative description of the e -invariant $K_3(\mathbb{C}_d) \rightarrow \mathbb{C}/\mathbb{Z}$. Abbreviate $X := \tau_{\geq 3}B\Gamma_\infty(\mathbb{C}_d)^+$ and consider the Bockstein exact sequence

$$(12) \quad H^3(X; \mathbb{C}) \longrightarrow H^3(X; \mathbb{C}/\mathbb{Z}) \xrightarrow{\delta} H^4(X; \mathbb{Z}) \longrightarrow H^4(X; \mathbb{C})$$

for the coefficients $\mathbb{Z} \rightarrow \mathbb{C} \rightarrow \mathbb{C}/\mathbb{Z}$.

By (8), the class $q_3^* T^* ch_2 = \frac{1}{2} q_3^* T^* s_2$ lies in $H^4(X; \mathbb{Z})$ and it maps to zero in $H^4(X; \mathbb{C})$ by Chern-Weil theory. Therefore there is a class $\epsilon \in H^3(X; \mathbb{C}/\mathbb{Z})$ such that $\delta \epsilon = \frac{1}{2} q_3^* T^* s_2$. If $x \in K_3(\mathbb{C}_d) = \pi_3(X)$, then $e(x) = \langle x^* \epsilon; [\mathbb{S}^3] \rangle$.

Let ψ_g be the composition $B\Gamma_g \rightarrow BSp_{2g}(\mathbb{Z}) \rightarrow BGL_{2g}(\mathbb{Z}) \rightarrow BGL_{2g}(\mathbb{C}_d)$ given by the action on the first cohomology of the surface. There is an induced map $\Psi : B\Gamma_\infty^+ \rightarrow B\Gamma_\infty(\mathbb{C}_d)^+$ in the stable range. The composition $T \circ \Psi$ differs from the map Φ described in Section 2. More precisely, let $c : BU \rightarrow BU$ be the map which sends a complex vector bundle V to $V \otimes_{\mathbb{R}} \mathbb{C} = V \oplus \bar{V}$. Then

$$(13) \quad T \circ \Psi \simeq c \circ \Phi$$

by the Hodge decomposition theorem.

Proposition 14. *The homomorphisms η and $e \circ (\Psi)_* : \pi_3(B\Gamma_\infty^+) \rightarrow \mathbb{C}/\mathbb{Z}$ coincide.*

Proof. Because we are only interested in π_3 , we may as well pass to the 2-connected covers of $B\Gamma_\infty^+$ and BU . We have to show that $(\tau_{\geq 3}\Psi)^*\epsilon = \bar{\zeta}_2$, where $\tau_{\geq 3}\Psi : \tau_{\geq 3}B\Gamma_\infty^+ \rightarrow \tau_3B\text{Gl}_\infty(\mathbb{C}_d)^+$ is induced from Ψ . Because $\bar{\zeta}_2$ is uniquely characterized by $\delta\bar{\zeta}_2 = q_3^*\zeta_2$, it suffices to show $\delta(\tau_{\geq 3}\Psi)^*\epsilon = q_3^*\zeta_2$. By the definition of ϵ and the functoriality of the connected covers, we obtain:

$$\delta(\tau_{\geq 3}\Psi)^*\epsilon = (\tau_{\geq 3}\Psi)^*\frac{1}{2}q_3^*T^*s_2 = (\tau_{\geq 3}\Psi)^*(\tau_{\geq 3}T)^*(\frac{1}{2}q_3^*s_2)$$

Using (8), it is easy to see that $(\tau_{\geq 3}c)^*q_3^*s_2 = 2s_2$. This relation, together with (13) and the definition of ζ_2 shows that

$$(\tau_{\geq 3}\Psi)^*(\tau_{\geq 3}T)^*(\frac{1}{2}q_3^*s_2) = (\tau_{\geq 3}\Phi)^*(q_3^*s_2) = q_3^*\zeta_2.$$

□

4. AN ACTION OF THE ICOSAHEDRAL GROUP ON A SURFACE

Consider a regular icosahedron \mathcal{I} in Euclidean 3-space, centered at 0 and such that all vertices lie on \mathbb{S}^2 . It has 20 faces (which are triangles), 12 vertices (at every vertex, exactly 5 edges and 5 faces meet) and 30 edges. Let $G \subset SO(3)$ be the symmetry group of the icosahedron. We choose a vertex, an edge and a face of the icosahedron. We denote the isotropy group of the midpoint of the edge by G_2 , the isotropy group of the midpoint of the face by G_3 and the isotropy group of the vertex by G_5 . These groups are cyclic of order 2, 3, 5, respectively, and we choose generators y_2, y_3, y_5 of these groups. The group G has order 60, it is generated by y_2, y_3 and y_5 and G is a perfect group.

Let \hat{G} the preimage of G under the 2-fold covering $\mathbb{S}^3 \rightarrow SO(3)$. The kernel of $\hat{G} \rightarrow G$ agrees with the center $Z(\hat{G})$ of \hat{G} . We denote the unique nontrivial element in $Z(\hat{G})$ by h . It is known that the extension $\mathbb{Z}/2 \rightarrow \hat{G} \rightarrow G$ is the universal central extension of G . The group \hat{G} is the *binary icosahedral group*.

We choose preimages x_i of the generators y_i . These set $\{x_2, x_3, x_5\}$ generates \hat{G} . The quotient $\hat{G} \backslash \mathbb{S}^3$ is the famous Poincaré homology 3-sphere.

Our analysis of G and \hat{G} begins with the description of the Sylow subgroups. Clearly $\langle x_2^2 \rangle$ and $\langle x_5^2 \rangle$ are the 3- and 5-Sylow subgroups of \hat{G} , respectively.

The quaternion group $Q_8 = \{\pm 1, \pm i, \pm j, \pm k\} \subset \mathbb{S}^3 \subset \mathbb{H}$ is a 2-Sylow subgroup of \hat{G} . The abelianization is $Q_8/\{\pm 1\} \cong V_4 := \mathbb{Z}/2 \times \mathbb{Z}/2$. Let U be the standard representation of $\hat{G} \subset \mathbb{S}^3 = \text{SU}(2)$ as a subgroup and denote the restriction to Q_8 by the same symbol. Let $u \in H^4(B\hat{G}; \mathbb{Z})$ be the second Chern class of U . Again, the restriction to BQ_8 will be denoted by u as well. Furthermore $\text{Hom}(Q_8; U(1)) \cong \text{Hom}(V_4; U(1)) \cong \mathbb{Z}/2 \times \mathbb{Z}/2$ (the last isomorphism

is not canonical). Let $a, b \in H^2(BQ_8; \mathbb{Z})$ be the first Chern classes of a pair of generators of $\text{Hom}(Q_8; U(1))$.

Proposition 15. *There is an isomorphism of rings $H^*(BQ_8; \mathbb{Z}) \cong \mathbb{Z}[a, b, u]/I$, where I is the ideal generated by $8u, 2a, 2b, a^2, b^2$ and $ab - 4u$.*

The cohomology ring $H^(B\hat{G}; \mathbb{Z})$ is isomorphic to $\mathbb{Z}[u]/(120u)$.*

Proof. By Poincaré duality, the cohomology of the closed oriented manifold \mathbb{S}^3/Q_8 is given by $H^*(\mathbb{S}^3/Q_8; \mathbb{Z}) \cong \mathbb{Z}$ for $* = 0, 3$, $H^2(\mathbb{S}^3/Q_8; \mathbb{Z}) \cong \mathbb{Z}/2 \times \mathbb{Z}/2$ and $H^*(\mathbb{S}^3/Q_8; \mathbb{Z}) = 0$ in all other cases. Now consider the commutative diagram of fibrations

$$(16) \quad \begin{array}{ccc} \mathbb{S}^3 & \longrightarrow & \mathbb{S}^3/Q_8 \\ \downarrow & & \downarrow \\ ES^3 & \longrightarrow & BQ_8 \\ \downarrow & & \downarrow \\ BS^3 & \xlongequal{\quad} & BS^3. \end{array}$$

The top horizontal map has degree 8. An application of the Leray-Serre spectral sequence shows that the additive structure of $H^*(BQ_8)$ is as asserted and that multiplication by u is an isomorphism $H^p(BQ_8) \rightarrow H^{p+4}(BQ_8)$ for all $p > 0$. The products on $H^2(BQ_8)$ are computed in [1, p. 60].

The computation for \hat{G} is similar, but easier, because the full structure follows from the spectral sequence. □

As a consequence of Proposition 15, the sum of the restriction maps to the Sylow subgroups $H^4(B\hat{G}) \rightarrow H^4(BQ_8) \oplus H^4(B\mathbb{Z}/3) \oplus H^4(B\mathbb{Z}/5) \cong \mathbb{Z}/8 \oplus \mathbb{Z}/3 \oplus \mathbb{Z}/5 \cong \mathbb{Z}/120$ is an isomorphism.

Remark 17. We will need a property of the representations of Q_8 . The representation U is (up to isomorphism) the only irreducible representation of Q_8 in which the central element h acts as -1 and not as the identity. It follows: If W is a Q_8 -representation, on which the central element acts by -1 , then W is a direct sum of copies of U .

Remark 18. The last fact we need is that any automorphism of the 3- and 5-Sylow subgroups of \hat{G} is induced by conjugation with an element in \hat{G} . The quickest way to see this uses the well-known isomorphism $\hat{G} \cong \text{Sl}_2(\mathbb{F}_5)$. This observation has the following consequence. If W is any representation of \hat{G} and $p = 3$ or $p = 5$, then the restriction of W to $\mathbb{Z}/p \subset \hat{G}$ is isomorphic to $n\mathbb{C} \oplus m(\bigoplus_{1 \leq r \leq p-1} L_{p,r})$ for appropriate $n, m \in \mathbb{N}$, where $L_{p,r}$ denotes the representation $1 \mapsto \exp(\frac{2\pi ir}{p})$ of \mathbb{Z}/p on \mathbb{C} .

Now we construct certain actions of \hat{G} on Riemann surfaces which in the end will give interesting elements in $\pi_3(B\Gamma_g^+)$. The construction is based on an easy lemma. Let $\mathcal{O}(k)$ be the k th tensor power of the Hopf bundle on $\mathbb{C}\mathbb{P}^1$.

The bundle $\mathcal{O}(k)$ is an $\mathrm{Sl}_2(\mathbb{C})$ -equivariant bundle over the $\mathrm{Sl}_2(\mathbb{C})$ -space \mathbb{CP}^1 . If k is odd, then the central element $-1 \in \mathrm{Sl}_2(\mathbb{C})$ acts as -1 on $\mathcal{O}(k)$ (i.e. on any fiber of the bundle). If k is even, then -1 acts trivially on $\mathcal{O}(k)$, i.e. the action descends to an action of $\mathbb{PSl}_2(\mathbb{C})$. Recall that the vector space of global holomorphic sections of $\mathcal{O}(k)$ can be identified with the vector space of homogeneous polynomials of degree k in $\mathbb{C}[t_1, t_2]$ (as $\mathrm{Sl}_2(\mathbb{C})$ -representations).

Lemma 19. *Let $G \subset \mathbb{PSl}_2(\mathbb{C})$ be a finite subgroup and let $\hat{G} \subset \mathrm{Sl}_2(\mathbb{C})$ be preimage of G in $\mathrm{Sl}_2(\mathbb{C})$ (it is a central extension of G by $\mathbb{Z}/2$). Let $m \in \mathbb{N}$ be positive. Let s be a G -invariant holomorphic section of $\mathcal{O}(2m)$ having only simple zeroes. Then there exists a connected Riemann surface F with a \hat{G} -action and a two-sheeted \hat{G} -equivariant branched covering $f : F \rightarrow \mathbb{CP}^1$, whose branch points are precisely the zeroes of s .*

If m is odd, then the central element $h \in \hat{G}$ acts as the hyperelliptic involution on F , if m is even, then h acts trivially on F .

Proof. Let $S \subset \mathcal{O}(2m)$ be the image of the section s . It is a surface of genus 0 and it is stable under the G -action on $\mathcal{O}(2m)$. Let $q : \mathcal{O}(m) \rightarrow \mathcal{O}(2m)$ be the squaring map, let $F := q^{-1}(S)$ and let $f := q|_F$. Clearly, F has a \hat{G} -action and f is equivariant. The other statements of the Lemma are trivial to verify. \square

From now on, let G again be the icosahedral group. The following example is the basis for the computation leading to Theorem 1.

Example 20. Let $z_1, \dots, z_{30} \in \mathbb{S}^2$ be the midpoints of the edges of the icosahedron and consider them as points on \mathbb{CP}^1 after the choice of a conformal map $\mathbb{S}^2 \cong \mathbb{CP}^1$. The precise value of the points does not play a significant role in this discussion.

Now we take holomorphic sections s_i , $i = 1, \dots, 30$, of the Hopf bundle $\mathcal{O}(1)$ with simple zeroes precisely at z_i . Such sections exist and are unique up to multiplication with a complex constant. Put $s := s_1 \otimes \dots \otimes s_{30} \in H^0(\mathbb{CP}^1, \mathcal{O}(30))$.

For $g \in \hat{G}$, there exists a $c(g) \in \mathbb{C}^\times$ with $gs = c(g)s$, because gs has the same zeroes as s . The map $c : g \mapsto c(g)$ is a homomorphism $\hat{G} \rightarrow \mathbb{C}^\times$. Since \hat{G} has no abelian quotient, c is constant. Thus s is a \hat{G} -invariant section.

If we apply the construction of Lemma 19 to s , we obtain a surface of genus 14 (by the Riemann-Hurwitz formula) with a \hat{G} -action.

Proposition 21. *The number of fixed points of the elements $x_2, x_3, x_5 \in \hat{G}$ in Example 20 is 2, 0, 0, respectively. All powers x_3^{2r} , $r \not\equiv 0 \pmod{3}$ and x_5^{2r} , $r \not\equiv 0 \pmod{5}$ have precisely 4 fixed points.*

The proof is an elementary consideration, using the fact that the number of fixed points of an automorphism of a Riemann surface is less or equal than $2g + 2$ with equality holding only for hyperelliptic involutions.

Of course, one can apply a similar construction to the midpoints of the faces or to the vertices of the icosahedron. The results are surfaces of lower genus and elements in $\pi_3(B\Gamma_g)^+$ of lower order.

5. COMPUTATIONS

The proof of Theorem 1 starts with the computation of the characteristic class ζ_2 for the bundle on $B\hat{G}$ given by the action $\rho : \hat{G} \rightarrow \Gamma_{14}$ constructed before.

Proposition 22. *The characteristic class $B\rho^*\zeta_2 \in H^4(B\hat{G}; \mathbb{Z}) \cong \mathbb{Z}/120$ has order 24.*

Proof. The \hat{G} -action on F is holomorphic by construction and therefore there is an induced linear action $\hat{G} \curvearrowright H^1(F; \mathcal{O})$. The cohomology class $B\rho^*\zeta_2$ of Proposition 22 is the same as the second Chern class of this linear \hat{G} -representation, which will be briefly denoted by c . According to the structure of $H^4(B\hat{G})$ described above, we need to show: The restriction of c to $H^4(B\mathbb{Z}/3)$ has order 3, the restriction to $H^4(\mathbb{Z}/5)$ is trivial and the restriction to $H^4(BQ_8)$ has order 8.

By the Dolbeault theorem, there is an isomorphism of complex \hat{G} -representations:

$$(23) \quad H^1(F; \mathcal{O}) \otimes_{\mathbb{R}} \mathbb{C} \cong H^1(F; \mathbb{C}).$$

The Lefschetz fixed point formula, applied to the result of Proposition 21, allows us to determine the decomposition of the representation of $\mathbb{Z}/3 \cong \hat{G}_{(3)} = \langle x_3^2 \rangle$ on the 28-dimensional space $H^1(F; \mathbb{C})$. The result is $H^1(F; \mathbb{C}) \cong 8\mathbb{C} \oplus 10L_1 \oplus 10L_2$. By (23) and Remark 18, it follows that $H^1(F; \mathcal{O}) \cong 4\mathbb{C} \oplus 5L_1 \oplus 5L_2$ as $\mathbb{Z}/3$ -modules. The Chern polynomial of L_n is $1 + nv$; $v \in H^2(\mathbb{Z}/3)$ an appropriate generator. It follows that $c|_{B\mathbb{Z}/3} = v^2$.

The result for the subgroup $\mathbb{Z}/5$ follows by an analogous argument. The decomposition of $H^1(F; \mathcal{O})$ as a $\mathbb{Z}/5$ -module is $2\mathbb{C} \oplus 3(L_1 \oplus L_2 \oplus L_3 \oplus L_4)$.

We have seen that the central element $h \in Q_8 \subset \hat{G}$ acts as a hyperelliptic involution. Thus it acts on $H^1(F; \mathcal{O})$ by -1 . By Remark 17, this implies that $H^1(F, \mathcal{O})$, as a Q_8 -module, decomposes into seven copies of the twodimensional representation U . The first Chern class of U is zero, while the second is a generator of $H^4(BQ_8)$. Thus $c|_{BQ_8}$ is seven times a generator and thus again a generator. □

Let $M = \mathbb{S}^3/\hat{G}$ be the Poincaré homology sphere and let $\theta_{M;\rho} \in \pi_3(B\Gamma_{\infty}^+)$ be the element determined by M and the action $\hat{G} \curvearrowright F_{14}$. In view of Theorem 2, the following proposition is enough to establish Theorem 1.

Proposition 24. *The order of $\eta(\theta) \in \mathbb{Q}/\mathbb{Z}$ is 24.*

Proof. Consider the 2-connected covering $q_3 : \tau_{\geq 3}B\Gamma_{\infty}^+ \rightarrow B\Gamma_{\infty}^+$ with homotopy fiber $K(\pi_2(B\Gamma_{\infty}^+); 1) \simeq \mathbb{S}^1$. By elementary obstruction theory (and the knowledge of the cohomology of $B\hat{G}$), there exists a unique (up to homotopy)

lift

(25)

$$\begin{array}{ccc}
 & & \tau_{\geq 3} B\Gamma_{\infty}^{+} \\
 & \nearrow \tilde{B}\rho & \downarrow \\
 B\hat{G}^{+} & \xrightarrow{B\rho} & B\Gamma_{\infty}^{+}.
 \end{array}$$

Comparing the Bockstein sequences for the spaces and maps $B\hat{G}^{+} \rightarrow \tau_{\geq 3} B\Gamma_{\infty}^{+} \rightarrow B\Gamma_{\infty}^{+}$, we see that $\tilde{B}\rho^{*} \bar{\zeta}_2 \in H^3(B\hat{G}^{+}; \mathbb{Q}/\mathbb{Z}) \cong \mathbb{Z}/120$ has order 24 (use (7), Lemma 3 and Proposition 22).

Now consider the map $\lambda^{+} : \mathbb{S}^3 \simeq M^{+} \rightarrow B\hat{G}^{+}$ which is induced from the classifying map $\lambda : M \rightarrow B\hat{G}$ of the universal covering of M . Observe that $\lambda^{*} : H^3(B\hat{G}^{+}; \mathbb{Q}/\mathbb{Z}) \rightarrow H^3(\mathbb{S}^3; \mathbb{Q}/\mathbb{Z})$ is injective, which follows from a straightforward consideration of the Leray-Serre spectral sequence for the fibration $\mathbb{S}^3 \rightarrow M \rightarrow B\hat{G}$.

Thus $\lambda^{*} \tilde{B}\rho^{*} \bar{\zeta}_2$ has order 24 in $H^3(\mathbb{S}^3; \mathbb{Q}/\mathbb{Z})$. But $\tilde{B}\rho \circ \lambda$ represents the element $\theta \in \pi_3(B\Gamma_{\infty}^{+})$ which we consider. \square

REFERENCES

- [1] M. F. Atiyah, Characters and cohomology of finite groups, *Inst. Hautes Études Sci. Publ. Math.* No. 9 (1961), 23–64. MR0148722 (26 #6228)
- [2] M. F. Atiyah and I. M. Singer, The index of elliptic operators. IV, *Ann. of Math.* (2) **93** (1971), 119–138. MR0279833 (43 #5554)
- [3] C. J. Earle and J. Eells, A fibre bundle description of Teichmüller theory, *J. Differential Geometry* **3** (1969), 19–43. MR0276999 (43 #2737a)
- [4] J. Ebert, Characteristic classes of spin surface bundles. Applications of the Madsen-Weiss theory, *Bonner Mathematische Schriften* 381, Bonn, 2006.
- [5] S. Galatius, Mod p homology of the stable mapping class group, *Topology* **43** (2004), no. 5, 1105–1132. MR2079997 (2006a:57020)
- [6] J. L. Harer, Stability of the homology of the mapping class groups of orientable surfaces, *Ann. of Math.* (2) **121** (1985), no. 2, 215–249. MR0786348 (87f:57009)
- [7] J. Harer, The second homology group of the mapping class group of an orientable surface, *Invent. Math.* **72** (1983), no. 2, 221–239. MR0700769 (84g:57006)
- [8] J. Harer, The third homology group of the moduli space of curves, *Duke Math. J.* **63** (1991), no. 1, 25–55. MR1106936 (92d:57012)
- [9] N. V. Ivanov, Complexes of curves and Teichmüller modular groups, *Uspekhi Mat. Nauk* **42** (1987), no. 3(255), 49–91, 255. MR0896878 (88g:32040)
- [10] N. V. Ivanov, On the homology stability for Teichmüller modular groups: closed surfaces and twisted coefficients, in *Mapping class groups and moduli spaces of Riemann surfaces (Göttingen, 1991/Seattle, WA, 1991)*, 149–194, *Contemp. Math.*, 150, Amer. Math. Soc., Providence, RI. MR1234264 (94h:57022)
- [11] J. D. S. Jones and B. W. Westbury, Algebraic K -theory, homology spheres, and the η -invariant, *Topology* **34** (1995), no. 4, 929–957. MR1362792 (98i:58217)
- [12] I. Madsen and U. Tillmann, The stable mapping class group and $Q(\mathbb{C}P_{\mp}^{\infty})$, *Invent. Math.* **145** (2001), no. 3, 509–544. MR1856399 (2002h:55011)
- [13] I. Madsen and M. Weiss, The stable moduli space of Riemann surfaces: Mumford’s conjecture, *Ann. of Math.* (2) **165** (2007), no. 3, 843–941. MR2335797 (2009b:14051)
- [14] J. Milnor, On the homology of Lie groups made discrete, *Comment. Math. Helv.* **58** (1983), no. 1, 72–85. MR0699007 (85b:57050)

- [15] S. Morita, Characteristic classes of surface bundles, *Invent. Math.* **90** (1987), no. 3, 551–577. MR0914849 (89e:57022)
- [16] J. Mukai, The S^1 -transfer map and homotopy groups of suspended complex projective spaces, *Math. J. Okayama Univ.* **24** (1982), no. 2, 179–200. MR0680191 (85d:55020)
- [17] J. Powell, Two theorems on the mapping class group of a surface, *Proc. Amer. Math. Soc.* **68** (1978), no. 3, 347–350. MR0494115 (58 #13045)
- [18] J. Rosenberg, *Algebraic K-theory and its applications*, Springer, New York, 1994. MR1282290 (95e:19001)
- [19] D. Quillen, Letter from Quillen to Milnor on $\text{Im}(\pi_i 0 \rightarrow \pi_i^s \rightarrow K_i \mathbf{Z})$, in *Algebraic K-theory (Proc. Conf., Northwestern Univ., Evanston, Ill., 1976)*, 182–188. Lecture Notes in Math., 551, Springer, Berlin. MR0482758 (58 #2811)
- [20] U. Tillmann, On the homotopy of the stable mapping class group, *Invent. Math.* **130** (1997), no. 2, 257–275. MR1474157 (99k:57036)

Received May 30, 2008; accepted Nov 6, 2008

Johannes Ebert

Rheinische Friedrich-Wilhelms-Universität Bonn, Mathematisches Institut
Endenicher Allee 60, 53115 Bonn, Germany

E-mail: ebert@math.uni-bonn.de

