


RESEARCH ARTICLE

Objectively measured versus self-reported physical activity in children and adolescents with cancer

Miriam Götte^{1*}, Corinna Caroline Seidel², Sabine Verena Kesting¹, Dieter Rosenbaum², Joachim Boos¹

1 Department of Pediatric Hematology and Oncology, University Children's Hospital Münster, Münster, Germany, **2** Movement Analysis Lab, Institute for Experimental Musculoskeletal Medicine, University Hospital Münster, Münster, Germany

* miriam.goette@ukmuenster.de



Abstract

Objective

Existing research recognizes low levels of physical activity in pediatric patients with cancer, but much uncertainty exists about their capability to self-reflect physical activity levels. The objective of this study was to compare results of subjective self-reports and objective accelerometers regarding levels of daily walking as well as moderate-to-vigorous physical activities.

Methods

Results of the objective assessment tool StepWatchTM Activity Monitor and self-reporting with a standardized questionnaire were compared in 28 children and adolescents during cancer treatment.

Results

The patients were 13.8±2.8 years of age and 3.4±2.0 months after cancer diagnosis. The Bland-Altman plots indicated a fairly symmetrical under- and over-estimation for daily minutes of walking with the limits of agreement ranging from -100.8 to 87.3 min (d = -6.7 min). Mean difference for moderate-to-vigorous physical activity was almost zero but limits of agreement are ranging from -126.8 to 126.9 min. The comparison for the days with at least 60 min of moderate-to-vigorous physical activity showed a marked difference with 3.0±2.6 self-reported days versus only 0.1±0.4 measured days.

Conclusions

These findings suggest that physical activity in pediatric cancer patients should preferably be assessed with objective methods. Greater efforts are needed to implement supervised exercise interventions during treatment incorporating methods to improve self-reflection of physical activity.

OPEN ACCESS

Citation: Götte M, Seidel CC, Kesting SV, Rosenbaum D, Boos J (2017) Objectively measured versus self-reported physical activity in children and adolescents with cancer. PLoS ONE 12(2): e0172216. doi:10.1371/journal.pone.0172216

Editor: Alejandro Lucía, Universidad Europea de Madrid, SPAIN

Received: October 10, 2016

Accepted: February 1, 2017

Published: February 16, 2017

Copyright: © 2017 Götte et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper.

Funding: This work was funded by the Cora-Lobscheid Stiftung <https://www.deutsches-stiftungszentrum.de/stiftungen/cora-lobscheid-stiftung> and the Kinderkrebshilfe Münster e.V. <http://www.kinderkrebshilfe-muenster.de/>.

Competing interests: The authors have declared that no competing interests exist.

Introduction

Adequate promotion of physical activities (PA) and sport is an essential prerequisite for motor development and enhances psychological and social health outcomes in children [1, 2]. Furthermore, PA and fitness during childhood are associated with numerous health benefits regarding physical (e.g. obesity) and psychological issues (e.g. depression) [3] and even with health conditions like metabolic syndrome during adulthood [4]. During cancer treatment, children and adolescents show considerably reduced levels of daily activities like walking and playing outdoors as well as significantly reduced minutes of exercise and sports per week [5]. This reduction of overall PA combined with the cancer disease and medical treatment lead to diminished motor performance and physical fitness at the end of acute cancer treatment [6, 7] and during aftercare [8, 9]. Valid tools to assess PA in this patient group are essential to define levels of inactivity and to measure the impact on activity levels in interventional studies including sports therapy. The current literature is lacking standardized methods to assess PA levels in pediatric oncology and studies either use subjective methods like questionnaires and activity scales [10, 5, 11, 12] or accelerometers [13–18]. Accelerometers are precise and valid but also complex, time-consuming and expensive and may not capture all movement whereas questionnaires and PA scales are less valid because they rely on the subject's recollection memory and interpretation [19, 20]. However, they allow to additionally integrate more in-depth questions about subjective perceptions and values regarding exercise as well as the precise type of exercise. Studies in healthy children and adolescents comparing subjective and objective measures of physical activity found acceptable correlations between self-reports and accelerometers for moderate physical activities but low correlations for vigorous activities with a tendency towards overestimating the active times [21, 22]. A search of the literature did not reveal any study that compared subjective and objective measures on PA in a childhood cancer population. The transparency of potential deviations between subjective self-reports and objective assessment is not only important for the design of future studies but for clinical practice because it reflects the capability of diseased children to self-estimate their levels of PA and therefore the capability to adhere to the recommendations and limits for being physically active during cancer treatment.

Aim of the present study was to compare results of subjective self-reports and objective accelerometers regarding levels of daily walking as well as moderate-to-vigorous physical activities (MVPA). We hypothesize that results of accelerometers and self-reports in children and adolescents with cancer will reveal significant deviations.

Methods

In the course of a previous study on physical activities, a sub-group of children and adolescents ($n = 29$) was gradually recruited for the comparison of subjective and objective assessment of PA levels and intensities during cancer treatment (Fig 1). All patients and parents gave informed written consent for participation. The study procedures were approved by the Ethics Committee of the General Medical Council Westfalen-Lippe and the Medical Faculty of Münster (file number 2012-035-f-S). Objective measures were obtained with the Step Watch 3™ Activity Monitor (SAM; Orthocare Innovations, Mountlake Terrace, WA 98043, USA). This sealed uniaxial accelerometer (7.5 x 5 x 2 cm; 38 g) counts the number of gait cycles (gcs) per time interval. Gait cycles are defined as two steps and the monitor was programmed to measure in 1-minute intervals. The device is attached to the ankle with an elastic strap. The SAM is an accurate and valid tool for measuring step activity and reportedly shows an accuracy of 99,87% in children [23]. It has been validated to provide accurate measurements in persons with slow gait [24] and severe disability [25]. Applicability in pediatric patients with leukemia,




Fig 1. Participant recruitment.

doi:10.1371/journal.pone.0172216.g001

lymphoma, brain tumor and bone tumor has been demonstrated [15, 18, 26]. Measurement outcomes of the SAM were the volume of activity per day (gcs per day) and intensity of activity (gcs per minute). In reference to the literature [27] 50 gcs/min (= 100 steps/min) were considered as threshold for moderate to vigorous activity levels and 20 gcs/min were defined as the threshold for continuous walking.

Subjective assessment of physical activities was conducted using the physical activity questionnaire (AQ) from the German Health Interview and Examination Survey for Children and Adolescents (KiGGS) of the Robert Koch Institute [28] with few additional cancer-specific

Table 1. Overview of the compared categories and respective measures of the AQ and SAM.

Category	SAM	AQ	Analysis
1: Walking (min/day)	Summarized daily minutes with > 20 gcs/min <i>min (mean)/day</i>	“How much do you walk (or go with crutches) during home stays?” <i>six answers options (from ‘never’ to ‘more than 2h/10km’ and more)</i>	Bland-Altman-Analysis, Spearman’s correlation
2: MVPA (days with ≥60 min/week)	Number of days with at least 60 minutes of >50 gcs/min <i>days/week</i>	“How many days per week are you physically active for at least 60 min during home stays?” <i>eight answer options (0 to 7 days)</i>	Mean ± SD
3: MVPA (min/week)	Summarized weekly minutes with > 50 gcs/min <i>min/week^a</i>	“Are you engaged in any sports at the moment or are you physically active in any kind? Which? How many minutes per week?” <i>type and minutes per week of up to four feasible sport activities</i>	Bland-Altman-Analysis, Spearman’s correlation

^a In study participants with less than seven days of wear time the hypothetical weekly score was calculated (daily mean of MVPA*7)

doi:10.1371/journal.pone.0172216.t001

questions. The original questionnaire has been tested for retest reliability with $r = 0.83$ and validity in comparison with the Sense Wear Pro2 was between $r = 0.56$ and $r = 0.66$ for leisure time sports and organized sports. The AQ consists of 45 items regarding physical activity levels before and during cancer treatment. Different questions assess daily PA (e.g. walking, playing outdoors) and moderate to vigorous physical activities (MVPA). A simple explanation of MVPA (elevated heart rate, be shortly put of breath and examples for activities) is given on the first page of the AQ. It took about 10–15 minutes to complete the questionnaire.

The comparison of SAM and AQ data was conducted for three categories: 1. Number of daily minutes of walking, 2. Number of days per week with at least 60min of MVPA and 3. Number of minutes with MVPA per week. Measurements by accelerometer (SAM) and self-report (AQ) are shown in Table 1. Participants filled in the AQ first as an unheralded self-reflection of physical activity levels without any previous motivational or educational talks. Thereafter they wore the SAM for seven consecutive days from morning after waking up until bedtime except during bathing or showering. All interested participants received an individual evaluation of their measured steps.

The accelerometer data was analyzed using the Step Watch 3™ Software 3.1 and raw data was exported into excel spread sheet for further analysis. Days with less than eight hours of wear time were excluded from the analysis. Commercial software (Graph Pad Prism, version 6.0) was used for statistical calculations and graphs. Bland-Altman plots were constructed to visualize the data and to assess agreement of AQ and SAM. To compare minutes of walking (category 1) of the questionnaire to the SAM data, the mean value of the marked time period (e.g. 22.5 min for marked answer 15–30 min) was used. Spearman’s analyses were computed to determine the degree of correlation between the two methods for minutes of walking and minutes of MVPA.

Results

Descriptive characteristics of the study population are presented in Table 2.

All 29 included children and adolescents returned the AQ and the SAM and there were no dropouts. However, one patient did not achieve the requested eight hours of wear time on any day and was therefore excluded from the analysis. Remaining 28 participants wore the SAM for 5.8 ± 2.8 days on average. Mean wearing time was 12.1 ± 1.5 hours per day. Objective measurement with the SAM revealed a mean of 52.0 ± 44.5 minutes of walking per day and 28.6 ± 31.3 minutes of MVPA per week (4.1 ± 4.5 min per day). Total number of gcs per day was 2859 ± 2028 gcs/day. Fig 2 shows the deviations of SAM (objective measure) and AQ

Table 2. Characteristics of the study population.

Characteristics	n (%)	Mean ± SD	Range
Age (years)	28 (100)	13.8 ± 2.8	8–20
BMI (kg/m ²)	28 (100)	20.2 ± 3.5	14.7–30.8
Gender			
Male	16 (57)		
Female	12 (43)		
Months since diagnosis		3.4 ± 2.0	1–9
Cancer type and age (years)			
Leukemia	13 (46)	12.9 ± 3.2	8–17
ALL	9		
AML	4		
Bone tumor	9 (32)	15.7 ± 2.1	13–20
Ewing sarcoma	3		
Osteosarcoma	6		
Localized at lower limb	3		
Localized at trunk/upper limb	6		
Lymphoma	2 (7)	13.0 ± 2.8	11–15
Other solid tumor	4 (14)	12.5 ± 1.3	11–14

Other solid tumor: soft tissue sarcoma (n = 2), brain tumor (n = 1), germ cell tumor (n = 1)

doi:10.1371/journal.pone.0172216.t002

(subjective measure) for the category “walking” (category 1). A value below zero indicates underestimation of walking by a patient whereas values above zero indicate overestimation. The mean difference (d) is -6.7 min and the limits of agreement are ranging from -100.8 to 87.3 min indicating a fairly equal under- and over-estimation of walking minutes per day. Differences between both tools are small among those walking <60 min and large beyond the




Fig 2. Bland-Altman plot of differences for daily walking minutes comparing AQ and SAM. x axis: mean of self-report/AQ and objective measure/SAM, y axis: differences between AQ and SAM, d = -6.7, 95% confidence limits.

doi:10.1371/journal.pone.0172216.g002




Fig 3. Bland-Altman plot of differences for weekly MVPA comparing AQ and SAM. x axis: mean of self-report/AQ and objective measure/SAM, y axis: differences between AQ and SAM, $d = 0.05$ min, 95% confidence limits.

doi:10.1371/journal.pone.0172216.g003

threshold of 60 min of walking. Due to the wide range and deviations in both directions (over- and underestimation) the absolute values of the differences were additionally calculated. Mean value of absolute deviations between AQ and SAM was 32.4 ± 35.5 min of walking per day. Spearman analysis indicates only a weak positive correlation between self-reported and objectively measured minutes of walking ($r = 0.3$; $P = 0.07$, graph not shown).

The comparison for the days with at least 60 min of MVPA (category 2) shows a marked difference with 3.0 ± 2.6 self-reported days (AQ) versus only 0.1 ± 0.4 measured days (SAM). Five patients (18%) reported to accumulate at least 60 min of MVPA on all days of the week, 11 patients reported three to six days with at least 60 min of MVPA, three patients reported one or two days and eight patient (29%) estimated that they did not meet this criterion on any day of the week. One patient forgot to answer this question. In contrast to this, the objective measurement of the SAM revealed that 26 of 28 patients (93%) did not accumulate 60 min of MVPA on any day of the week. One participant each was physically active with moderate to vigorous intensity on one and two days of the week.

Fig 3 compares minutes of MVPA per week as measured by SAM and AQ (category 3). The diagram shows that the mean difference (d) for MVPA measured by AQ and SAM is almost zero (0.05 min) but the limits of agreement are ranging from -126.8 to 126.9 min. Differences are small at low levels (<30–40 min) with discrepancies occurring at > 50 min MVPA. Mean value of absolute deviations between AQ and SAM was 42.1 ± 49.9 minutes of MVPA per week. Spearman analysis indicates that there is only a weak correlation between self-reported and objectively measured weekly minutes of MVPA ($r = 0.1$; $P = 0.12$, graph not shown).

Table 3 demonstrates the differences of self-estimation between patients with hematological tumors and patients with solid tumors. Correct estimation was defined as a relative deviation between self-reported and objective measure below 30% and over- and underestimation as deviations over $\pm 30\%$, respectively. Almost 50% of the children with leukemia or lymphoma

Table 3. Self-estimation of physical activity differentiated for patients with hematological vs. solid tumors.

	All patients, n (%)	Leukemia/ lymphoma patients, n (%)	Patients with solid Tumors, n (%)
Daily walking			
correct estimation	7 (25.0)	3 (20.0)	4 (30.8)
overestimation	10 (35.7)	7 (46.7)	3 (23.1)
underestimation	11 (39.3)	5 (33.3)	6 (46.2)
Weekly MVPA			
correct estimation	6 (23.1)	4 (30.8)	2 (15.4)
overestimation	6 (23.1)	3 (23.1)	3 (23.1)
underestimation	14 (53.8)	6 (46.2)	8 (61.5)

doi:10.1371/journal.pone.0172216.t003

overestimated their daily minutes of walking whereas MVPA was more often underestimated. Patients with solid tumors, in this sample particularly bone tumor patients, tended to underestimate their levels of daily walking as well as levels of MVPA.

Discussion

The present study used a standardized questionnaire as a subjective self-report and the Step Watch 3™ Activity Monitor as an objective tool to compare minutes of walking as well as minutes of MVPA. Our hypothesis that results from objective and subjective measurements of physical activity deviate from each other in children and adolescents undergoing cancer treatment was confirmed by the present data. Although mean differences between both tools seem to be negligible at first (6.7 min for walking and 0 min for MVPA), further analysis revealed large individual deviations that are just evened out in the mean and indicates almost equal over- and underestimation of activity levels. The results support our hypothesis that young patients with cancer have problems to reliably self-report levels of PA and deviations are increasing with amount of PA. Problems regarding self-reflection of physical activities have been reported in children with other chronic diseases like juvenile dermatomyositis and juvenile systemic lupus erythematosus [29] and the presented study reveals similar results for pediatric patients with cancer. In comparison with healthy children, the direction of self-reflection is uncommon in our study, because children typically overestimate their physical activity levels in self-reports [30, 22]. A possible explanation for poor correlations in the present study may be the changed daily routine for children during cancer treatment. Children are isolated from school and often lose contact to their social surroundings and activities. This loss of regular activity times like physical education and engagement in sport clubs presumably makes it very difficult for them to remember and categorize moments of PA. The overall low levels of PA during treatment may influence self-reports in a way that children refer their answers to their desire for being physically active. The unexpected underestimation of PA and MVPA in a relevant number of children in this study may reflect their own exaggerated feelings of being inactive due to limitations in body function and overall weakness [31]. This phenomenon seems to be especially present in patients with bone tumors and other solid tumors. More than 60% of these children underestimated their weekly minutes of MVPA by more than 30%, i.e. the accelerometer measured more minutes of MVPA than the children assumed. To support the patients' self-confidence it might be beneficial for the children and adolescents to receive individual feedback about their daily activities by using accelerometers. This could possibly influence their own perception of abilities and physical activity in a positive way and demonstrate that even moderate to vigorous levels of PA are feasible.

Some limitations should be acknowledged regarding the study design. This study included a limited number of participants; however, with 28 children and different tumor entities a realistic insight into the topic should be possible. Furthermore, the marginal deviations (10%) regarding overall physical activity levels between the patients of this study and a comparison group from a previous study [18] indicate that the children of this study seem to be a representative cohort for pediatric cancer patients during treatment. Secondly we cannot specify the percentage of cancer treatment influencing the results in addition to the general problems in self-estimating physical activities in children. Because AQ showed good reliability and validity in previous studies with healthy children [32] it can be assumed that differences are mainly due to a loss of self-estimation capacities in children during the phase of cancer treatment. It has to be noticed that some participants have not achieved the requested seven days of SAM wear time; however according to a previous study a duration of four days of monitoring has been described as sufficient to measure habitual physical activity levels when using sealed pedometers in children [33]. To our knowledge this is the first study to assess self-reflection of physical activity levels in a cohort of pediatric patients with cancer using objective accelerometry and self-reports. There is no control group of healthy children as it was not the purpose of this study to compare self-reflection between diseased and healthy children. Therefore, the inclusion of such a control group would not have added necessary information to this study. All methods used have been tested for validity in previous studies [23–25] and for applicability in children during cancer treatment [5, 15, 18, 26].

In conclusion, objective measures should be preferentially used for the assessment of physical activities in children and adolescents with cancer to ensure accurate and reliable data. Self-reports may complementarily assess categories of activities and sports not including steps as well as expectations and attitudes towards exercise and physical activities. Findings of this study may also have important practical implications for children and adolescents with cancer. Exercise interventions during treatment should be supervised to ensure controlled and safe conditions. These interventions may also be guided with objective tools. Secondly, these interventions should incorporate methods to improve self-reflection of physical activity levels and intensity and overall body awareness, e.g. through activity monitors with biofeedback. Further research with larger samples of children and adolescents with cancer should be undertaken to investigate in detail the influences of tumor entities, age, gender and treatment phase on self-reflection of physical activity.

Author Contributions

Conceptualization: MG CCS JB.

Data curation: MG.

Formal analysis: MG.

Funding acquisition: JB.

Investigation: MG SVK.

Methodology: MG CCS DR.

Project administration: MG SVK JB.

Visualization: MG.

Writing – original draft: MG.

Writing – review & editing: MG CCS SVK DR JB.

References

1. Eime RM, Young JA, Harvey JT, Charity MJ, Payne WR. A systematic review of the psychological and social benefits of participation in sport for children and adolescents: informing development of a conceptual model of health through sport. *Int J Behav Nutr Phys Act.* 2013; 10: 98. doi: [10.1186/1479-5868-10-98](https://doi.org/10.1186/1479-5868-10-98) PMID: [23945179](https://pubmed.ncbi.nlm.nih.gov/23945179/)
2. Timmons BW, Naylor P-J, Pfeiffer KA. Physical activity for preschool children-how much and how? *Can J Public Health.* 2007; 98: 122–134.
3. Janssen I, Leblanc AG. Systematic review of the health benefits of physical activity and fitness in school-aged children and youth. *Int J Behav Nutr Phys Act.* 2010; 7: 40. doi: [10.1186/1479-5868-7-40](https://doi.org/10.1186/1479-5868-7-40) PMID: [20459784](https://pubmed.ncbi.nlm.nih.gov/20459784/)
4. Fraser BJ, Huynh QL, Schmidt MD, Dwyer T, Venn AJ, Magnussen CG. Childhood Muscular Fitness Phenotypes and Adult Metabolic Syndrome. *Med Sci Sports Exerc.* 2016; 48: 1715–1722. doi: [10.1249/MSS.0000000000000955](https://doi.org/10.1249/MSS.0000000000000955) PMID: [27128670](https://pubmed.ncbi.nlm.nih.gov/27128670/)
5. Götte M, Kesting S, Winter C, Rosenbaum D, Boos J. Comparison of self-reported physical activity in children and adolescents before and during cancer treatment. *Pediatr Blood Cancer.* 2014; 61: 1023–1028. doi: [10.1002/pbc.24898](https://doi.org/10.1002/pbc.24898) PMID: [24357259](https://pubmed.ncbi.nlm.nih.gov/24357259/)
6. Götte M, Kesting S V, Winter CC, Rosenbaum D, Boos J. Motor performance in children and adolescents with cancer at the end of acute treatment phase. *Eur J Pediatr.* 2015; 174: 791–799. doi: [10.1007/s00431-014-2460-x](https://doi.org/10.1007/s00431-014-2460-x) PMID: [25428233](https://pubmed.ncbi.nlm.nih.gov/25428233/)
7. San Juan AF, Chamorro-Viña C, Maté-Muñoz JL, Fernández del Valle M, Cardona C, Hernández M, et al. Functional capacity of children with leukemia. *Int J Sports Med.* 2008; 29: 163–167. doi: [10.1055/s-2007-964908](https://doi.org/10.1055/s-2007-964908) PMID: [17879894](https://pubmed.ncbi.nlm.nih.gov/17879894/)
8. Beulertz J, Bloch W, Prokop A, Baumann FT. Specific deficit analyses in motor performance and quality of life of pediatric cancer patients—a cross-sectional pilot study. *Pediatr Hematol Oncol.* 2013; 30: 336–347. doi: [10.3109/08880018.2013.776155](https://doi.org/10.3109/08880018.2013.776155) PMID: [23495850](https://pubmed.ncbi.nlm.nih.gov/23495850/)
9. Kesting SV, Götte M, Seidel CC, Rosenbaum D, Boos J. Motor Performance After Treatment for Pediatric Bone Tumors. *J Pediatr Hematol Oncol* 2015; 37: 509–514. doi: [10.1097/MPH.0000000000000396](https://doi.org/10.1097/MPH.0000000000000396) PMID: [26207777](https://pubmed.ncbi.nlm.nih.gov/26207777/)
10. Finnegan L, Wilkie DJ, Wilbur J, Campbell RT, Zong S, Katula S. Correlates of physical activity in young adult survivors of childhood cancers. *Oncol Nurs Forum.* 2007; 34: E60. doi: [10.1188/07.ONF.E60-E69](https://doi.org/10.1188/07.ONF.E60-E69) PMID: [17878118](https://pubmed.ncbi.nlm.nih.gov/17878118/)
11. Kesting S, Götte M, Winter C, Rosenbaum D, Boos J. One in Four Questioned Children Faces Problems Regarding Reintegration Into Physical Education at School After Treatment for Pediatric Cancer. *Pediatr Blood Cancer.* 2016; 63: 737–739. doi: [10.1002/pbc.25852](https://doi.org/10.1002/pbc.25852) PMID: [26681662](https://pubmed.ncbi.nlm.nih.gov/26681662/)
12. Rueegg CS, Gianinazzi ME, Michel G, von der Weid NX, Bergstraesser E, Kuehni CE. Do childhood cancer survivors with physical performance limitations reach healthy activity levels? *Pediatr Blood Cancer.* 2013; 60: 1714–1720. doi: [10.1002/pbc.24595](https://doi.org/10.1002/pbc.24595) PMID: [23681516](https://pubmed.ncbi.nlm.nih.gov/23681516/)
13. Aznar S, Webster AL, San Juan AF, Chamorro-Viña C, Maté-Muñoz JL, Moral S, et al. Physical activity during treatment in children with leukemia: a pilot study. *Appl Physiol Nutr Metab.* 2006; 31: 407–413. doi: [10.1139/h06-014](https://doi.org/10.1139/h06-014) PMID: [16900230](https://pubmed.ncbi.nlm.nih.gov/16900230/)
14. Braam KI, van Dijk-Lokkart EM, Kaspers GJ, Takken T, Huisman J, Bierings MB, et al. Cardiorespiratory fitness and physical activity in children with cancer. *Support Care Cancer.* 2016; 24: 2259–2268. doi: [10.1007/s00520-015-2993-1](https://doi.org/10.1007/s00520-015-2993-1) PMID: [26581899](https://pubmed.ncbi.nlm.nih.gov/26581899/)
15. Müller C, Winter C, Boos J, Gosheger G, Harges J, Vieth V, et al. Effects of an exercise intervention on bone mass in pediatric bone tumor patients. *Int J Sports Med.* 2014; 35: 696–703. doi: [10.1055/s-0033-1358475](https://doi.org/10.1055/s-0033-1358475) PMID: [24408763](https://pubmed.ncbi.nlm.nih.gov/24408763/)
16. Orsey AD, Wakefield DB, Cloutier MM. Physical Activity (PA) and Sleep Among Children and Adolescents With Cancer. *Pediatr Blood Cancer* 2013; 60: 1908–1913. doi: [10.1002/pbc.24641](https://doi.org/10.1002/pbc.24641) PMID: [23775811](https://pubmed.ncbi.nlm.nih.gov/23775811/)
17. Tan SY, Poh BK, Chong HX, Ismail MN, Rahman J, Zarina AL, et al. Physical activity of pediatric patients with acute leukemia undergoing induction or consolidation chemotherapy. *Leuk Res.* 2013; 37: 14–20. doi: [10.1016/j.leukres.2012.09.005](https://doi.org/10.1016/j.leukres.2012.09.005) PMID: [23099236](https://pubmed.ncbi.nlm.nih.gov/23099236/)
18. Winter C, Müller C, Brandes M, Brinkmann A, Hoffmann C, Harges J, et al. Level of activity in children undergoing cancer treatment. *Pediatr Blood Cancer.* 2009; 53: 438–443. doi: [10.1002/pbc.22055](https://doi.org/10.1002/pbc.22055) PMID: [19415742](https://pubmed.ncbi.nlm.nih.gov/19415742/)
19. Shephard RJ. Limits to the measurement of habitual physical activity by questionnaires. *Br J Sports Med.* 2003; 37: 197–206. doi: [10.1136/bjism.37.3.197](https://doi.org/10.1136/bjism.37.3.197) PMID: [12782543](https://pubmed.ncbi.nlm.nih.gov/12782543/)

20. Vanhees L, Lefevre J, Philippaerts R, Martens M, Huygens W, Troosters T, et al. How to assess physical activity? How to assess physical fitness? *Eur J Cardiovasc Prev Rehabil.* 2005; 12: 102–14. PMID: [15785295](#)
21. Manios Y, Androutsos O, Moschonis G, Birbilis M, Maragkopoulou K, Giannopoulou A, et al. Criterion validity of the physical activity questionnaire for schoolchildren (PAQ-S) in assessing physical activity levels: the healthy growth study. *J Sports Med Phys Fitness.* 2013; 53: 502–508 PMID: [23903530](#)
22. LeBlanc AG, Janssen I. Differences between self-reported and accelerometer measured moderate-to-vigorous physical activity in youth. *Pediatr Exerc Sci.* 2010; 22: 523–534. PMID: [21242602](#)
23. McDonald CM, Widman L, Abresch RT, Walsh SA, Walsh DD. Utility of a step activity monitor for the measurement of daily ambulatory activity in children. *Arch Phys Med Rehabil.* 2005; 86: 793–801. doi: [10.1016/j.apmr.2004.10.011](#) PMID: [15827934](#)
24. Storti KL, Pettee KK, Brach JS, Talkowski JB, Richardson CR, Kriska AM. Gait speed and step-count monitor accuracy in community-dwelling older adults. *Med Sci Sports Exerc.* 2008; 40: 59–64. doi: [10.1249/mss.0b013e318158b504](#) PMID: [18091020](#)
25. Sandroff BM, Motl RW, Pilutti LA, Learmonth YC, Ensari I, Dlugonski D, et al. Accuracy of StepWatch and ActiGraph accelerometers for measuring steps taken among persons with multiple sclerosis. *PLoS One.* 2014; 9: e93511. doi: [10.1371/journal.pone.0093511](#) PMID: [24714028](#)
26. Winter CC, Müller C, Harges J, Gosheger G, Boos J, Rosenbaum D. The effect of individualized exercise interventions during treatment in pediatric patients with a malignant bone tumor. *Support Care Cancer.* 2013; 21: 1629–1636. doi: [10.1007/s00520-012-1707-1](#) PMID: [23292667](#)
27. Tudor-Locke C, Hatano Y, Pangrazi RP, Kang M. Revisiting “how many steps are enough?”. *Med Sci Sports Exerc.* 2008; 40: 537–543.
28. Bös K, Worth A, Opper E, Oberger J, Woll A. Motorik-Modul: Eine Studie zur motorischen Leistungsfähigkeit und körperlich-sportlichen Aktivität von Kindern und Jugendlichen in Deutschland. The Motorik-Modul: physical fitness and physical activity in German children and adolescents. 1st ed. Baden-Baden: Nomos Verlag; 2009.
29. Pinto AJ, Roschel H, Benatti FB, de Sá Pinto AL, Sallum AME, Silva CA, et al. Poor agreement of objectively measured and self-reported physical activity in juvenile dermatomyositis and juvenile systemic lupus erythematosus. *Clin Rheumatol.* 2016; 35: 1507–1514. doi: [10.1007/s10067-016-3234-9](#) PMID: [27021334](#)
30. Welk GJ, Corbin CB, Dale D. Measurement Issues in the Assessment of Physical Activity in Children. *Res Q Exerc Sport.* 2000; 71: 59–73. doi: [10.1080/02701367.2000.11082788](#) PMID: [25680015](#)
31. Darcy L, Björk M, Knutsson S, Granlund M, Enskär K. Following Young Children’s Health and Functioning in Everyday Life Through Their Cancer Trajectory. *J Pediatr Oncol Nurs.* 2015; 33: 173–189. doi: [10.1177/1043454215610489](#) PMID: [26655332](#)
32. Woll A, Kurth B-M, Opper E, Worth A, Bös K. The “Motorik-Modul” (MoMo): physical fitness and physical activity in German children and adolescents. *Eur J Pediatr.* 2011; 170: 1129–1142. doi: [10.1007/s00431-010-1391-4](#) PMID: [21318230](#)
33. Ozdoba R, Corbin C, Le Masurier G. Does Reactivity Exist in Children When Measuring Activity Levels With Unsealed Pedometers? *Pediatr Exerc Sci.* 2004; 16: 158–166.